

010307
0400533

010307
0400533

Drug Facts

Active ingredient
(In each 5 mL teaspoonful)
Dextromethorphan polistirex, equivalent to 50 mg dextromethorphan hydrobromide USP

Purpose
Cough suppressant

Use
temporarily relieves cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, Parkinson's disease, and high blood pressure) or if you have taken a MAOI in the last 14 days. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

■ chronic cough that lasts as long as 7 days, cough comes back, or occurs with fever, rash or headache that lasts, these could be signs of a serious condition.
■ cough that occurs with too much phlegm (mucus).
■ Stop use and ask a doctor if cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts, these could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Shake bottle well before use
■ Shake as follows or as directed by a doctor
■ adults and children 12 years of age and over 2 teaspoonfuls every 12 hours, not to exceed 4 teaspoonfuls in 24 hours
■ children 6 to under 12 years of age 1 teaspoonful every 12 hours, not to exceed 2 teaspoonfuls in 24 hours
■ children 2 to under 6 years of age 1/2 teaspoonful every 12 hours, not to exceed 1 teaspoonful in 24 hours
■ children under 2 years of age consult a doctor

Other information

■ each 5 mL teaspoonful contains: sodium 8 mg
■ store at 20°-25°C (68°-77°F)

Inactive ingredients citric acid anhydrous, D&G Red #33, edetate disodium, ethylcellulose, FD&C Blue #1, flavor, high fructose corn syrup, menthylparaben, polyethylene glycol 550, polysorbate 80, purified water, propylparaben, purified water, saccharin, vegetable oil, vegetable oil, xanthan gum

Questions? 1-866-682-4639

© 2007, distributed by Adams Respiratory Therapeutics, Fort Worth, TX 76155

12 Hour Cough Relief
Delsym

NDC 63824-171-65

12 Hour Cough Relief
Delsym

dextromethorphan polistirex
extended-release suspension

**COUGH
SUPPRESSANT**

NEW GRAPE
FLAVOR



Grape-Flavored Liquid



Alcohol-free
148 mL (5 fl oz)

3-63824-171-65-4

Lot No.
Exp. Date

12 Hour Cough Relief
Delsym

dextromethorphan polistirex
extended-release suspension

**COUGH
SUPPRESSANT**

DELSYM® DOSING
SHAKE WELL BEFORE USE.


Age (yr)	Dose (1 TSP = 5 mL)
12 years to adult	2 TEASPOONFULS EVERY 12 HOURS
6 to under 12	1 TEASPOONFUL EVERY 12 HOURS
2 to under 6	1/2 TEASPOONFUL EVERY 12 HOURS

See back panel for full dosing directions.

Please visit our Web site:
www.delsym.com

Contains
No Fever Reducer
or Pain Reliever
Great Grape Taste

FILE INFORMATION
Job Description:
Drawing #: #002
Carton Size (Buil
Software: Adobe
Date Work Perfor
Colors Used: 4/C
Fonts Used: Fruit
Helv
Additional Inform



FILE INFORMATION

Job Description: Delsym Adult Grape 5-oz. Label

Drawing #: Not Available

Label Size: 6-5/16"w x 2-1/2"h

Software: Adobe Illustrator 12

Date Work Performed: 02-13-07

Colors Used: PMS 527 + PMS 2738 + Process Yellow

Fonts Used: Frutiger, Helvetica Condensed,
Helvetica Neue Cond, Helvetica Neue Ext

Additional Information:

glue area
no ink & coating

Drug Facts

Active ingredient (in each 5 mL teaspoonful) Purpose
Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide, USP. Cough suppressant

Use Temporarily relieves cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- chronic cough that lasts as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Shake bottle well before use

■ dose as follows or as directed by a doctor	
adults and children 12 years of age and over	2 teaspoonfuls every 12 hours, not to exceed 4 teaspoonfuls in 24 hours
children 6 to under 12 years of age	1 teaspoonful every 12 hours, not to exceed 2 teaspoonfuls in 24 hours
children 2 to under 6 years of age	1/2 teaspoonful every 12 hours, not to exceed 1 teaspoonful in 24 hours
children under 2 years of age	consult a doctor

Other information

■ each 5 mL teaspoonful contains: sodium 6 mg

■ store at 20°-25°C (68°-77°F)

Inactive ingredients: citric acid anhydrous, D&C Red #33, edetate disodium, ethylcellulose, FD&C Blue #1, flavor high fructose corn syrup, methylparaben, polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, sorbic acid, vegetable oil, xanthan gum

Questions? 1-866-682-4639



ADAMS
RESPIRATORY
THERAPEUTICS

010307
4000529

© 2007. Distributed by Adams Respiratory Therapeutics, Fort Worth, TX 76155

GLUE AREA
NO INK & COATING

12 Hour Cough Relief Delsym

NDC 63824-171-63

Delsym

dextromethorphan polistirex
extended-release suspension
**COUGH
SUPPRESSANT**

**NEW GRAPE
FLAVOR**



Grape-Flavored Liquid



Alcohol-free
89 mL (3 fl oz)

**Also Available
In Great
Orange Flavor**



**Contains
No Fever Reducer
or Pain Reliever
Great Grape Taste**

12 Hour Cough Relief Delsym

dextromethorphan polistirex
extended-release suspension
**COUGH
SUPPRESSANT**

DELSYM® DOSING SHAKE WELL BEFORE USE.

Age (yr)	Dose (1 TSP = 5 mL)
12 years to adult	2 TEASPOONFULS EVERY 12 HOURS
6 to under 12	1 TEASPOONFUL EVERY 12 HOURS
2 to under 6	1/2 TEASPOONFUL EVERY 12 HOURS

See back panel for full dosing directions.

Please visit our Web site:
www.delsym.com

**Contains
No Fever Reducer
or Pain Reliever
Great Grape Taste**

GLUE AREA

NO INK & COATING

FILE INFORMATION
Job Description
Drawing #: #C
Carton Size (B
Software: Ado
Date Work Per
Colors Used: 4
Fonts Used: F
Additional Info



2 1/8"

5 5/8"

FILE INFORMATION

Job Description: Delsym Adult Grape 3-oz. Label

Drawing #: Not Available

Label Size: 5-5/8"w x 2-1/8"h

Software: Adobe Illustrator 12

Date Work Performed: 02-13-07

Colors Used: PMS 527 + PMS 2738 + Process Yellow

Fonts Used: Frutiger, Helvetica Condensed,
Helvetica Neue Cond, Helvetica Neue Ext

Additional Information:

FILE INFORMATION

Job Description: Delsym
Drawing #: #C020280B
Carton Size (Built): 2-15
Software: Adobe Illustrat
Date Work Performed: 0
Colors Used: 4/C + PMS
Fonts Used: Frutiger, Hel
Helv Neue C
Additional Information:

12 Hour Cough Relief Delsym

NDC 63824-172-65

12 Hour Cough Relief
Delsym
dextromethorphan polistirex
extended-release suspension
**COUGH
SUPPRESSANT**

12 Hour Cough Relief
Delsym
dextromethorphan polistirex
extended-release suspension
**COUGH
SUPPRESSANT**

12 Hour Cough Relief
Delsym
dextromethorphan polistirex
extended-release suspension
**COUGH
SUPPRESSANT**

**NEW GRAPE
FLAVOR**



DELSYM® DOSING

SHAKE WELL BEFORE USE.

Age (yr)	Dose (1 TSP = 5 mL)
Under 2	CONSULT DOCTOR
2 to under 6	1/2 TEASPOONFUL EVERY 12 HOURS
6 to under 12	1 TEASPOONFUL EVERY 12 HOURS

See back panel for full dosing directions.

Please visit our Web site:
www.delsym.com

**Contains
No Fever Reducer
or Pain Reliever**
Great Grape Taste



Alcohol-free
148 mL (5 fl oz)

Grape-Flavored Liquid
for Children & Adults

3-63824-172-65-1

Lot No.:
Exp. Date:

Delsym

guaranteed to work or is ☒ with genuine
pharmaceutical quality. ☐ ask for 00
TAMPER EVIDENT SEAL

Drug Facts

Active Ingredient

(In each 5 mL teaspoonful)

Dextromethorphan polistirex equivalent to
30 mg dextromethorphan hydrobromide. Cough suppressant
as may occur with the common cold or influenza viruses.

Purpose

Use temporarily relieves cough due to minor throat and bronchial irritation

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor
(MAOI) (certain drugs for depression, psychiatric or emotional conditions, or
Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do
not know if your prescription drug contains an MAOI, ask a doctor or
pharmacist before taking this product.

Ask a doctor before use if you have

■ chronic cough that lasts as long as 3 weeks, asthma or emphysema
■ cough that occurs with or without fever, chest pain, or more than 7 days' cough comes
back, or cough that is worse at night

Stop use and ask a doctor if cough persists more than 7 days' cough comes
back, or cough that is worse at night, fever, chest or headache that lasts. These could be signs
of a more serious condition.

Use as directed. Do not use for more than 7 days. If symptoms do not improve, ask a
health professional before use.
Keep out of reach of children. In case of overdose, get medical help or
contact a Poison Control Center right away.

Directions

Shake bottle well before use

■ Use as follows or as directed by a doctor

adults and children 12 years of age and over	2 teaspoonfuls every 12 hours, not to exceed 4 teaspoonfuls in 24 hours
children 6 to under 12 years of age	1 teaspoonful every 12 hours in 24 hours, not to exceed 2 teaspoonfuls in 24 hours
children 2 to under 6 years of age	1/2 teaspoonful every 12 hours, not to exceed 1 teaspoonful in 24 hours
children under 2 years of age	Consult a doctor

Other Information

■ each 5 mL teaspoonful contains sodium 8 mg
■ store at 20°-25°C (68°-77°F)

Inactive ingredients: citric acid anhydrous, D&C Red #28, edetate
disodium, ethylcellulose, FD&C Blue #1, flavor, high fructose corn syrup,
methylparaben, polyethylene glycol 3350, polyethylene glycol,
propylparaben, purified water, sucrose, triacetin, vegetable oil, xanthan gum

Questions? 1-866-682-4639

**ADAM'S
RESPIRATORY
THERAPEUTICS**

021307
4000534

© 2007. Distributed by Adams Respiratory Therapeutics, Fort Worth, TX 76155

NDC 63824-172-65

Delsym

dextromethorphan polistirex
extended-release suspension

COUGH SUPPRESSANT

12 Hour Cough Relief

Contains No Fever Reducer
or Pain Reliever

Alcohol-free

Grape-Flavored Liquid
for Children & Adults

148 mL (5 fl oz)

© 2007, Distributed by
Adams Respiratory Therapeutics
Fort Worth, TX 76155

Lot:
Exp:

021207 4000522

INDICATIONS: Temporarily relieves cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants.

DIRECTIONS: SHAKE BOTTLE WELL BEFORE USING.

Dose as follows or as directed by a doctor:

Adults and Children 12 years of age and over: 2 teaspoonsful every 12 hours, not to exceed 4 teaspoonsful in 24 hours.

Children 6 to under 12 years of age: 1 teaspoonful every 12 hours, not to exceed 2 teaspoonsful in 24 hours.

Children 2 to under 6 years of age: 1/2 teaspoonful every 12 hours, not to exceed 1 teaspoonful in 24 hours.

Children under 2 years of age: Consult a doctor.

WARNINGS: Do not take this product for chronic cough that lasts as occurs with smoking, asthma, or emphysema, or if cough occurs with too much phlegm (mucus) unless directed by a doctor. If cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts, consult a doctor. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

DRUG INTERACTION PRECAUTION: Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

ACTIVE INGREDIENT: Each teaspoonful (5 mL) contains dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide.

TAMPER EVIDENT: Do not use if the neckband printed with is broken or missing.

Store at 20°-25°C (68°-77°F).

Questions? 1-866-682-4639

3 63824 17265 1

FILE INFORMATION

Job Description: Delsym Pediatric Grape 5-oz. Label

Drawing #: Not Available

Label Size: 6-5/16"w x 2-1/2"h

Software: Adobe Illustrator 12

Date Work Performed: 02-13-07

Colors Used: PMS 527 + PMS 2738 + Process Yellow

Fonts Used: Frutiger, Helvetica Condensed,
Helvetica Neue Cond, Helvetica Neue Ext

Additional Information:

•guissus is or unorg is 2 with printed and primed with and
If asen you Og :JNEIDIVIA HERJIMV

Drug Facts

Active ingredient (in each 5 mL teaspoonful) **Purpose**
Dextromethorphan polistirex equivalent to
30 mg dextromethorphan hydrobromide..... Cough suppressant

Use temporarily relieves cough due to minor throat and bronchial
irritation as may occur with the common cold or inhaled irritants

Warnings
Do not use if you are now taking a prescription monoamine oxidase
inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional
conditions, or Parkinson's disease), or for 2 weeks after stopping the
MAOI drug. If you do not know if your prescription drug contains an
MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
■ chronic cough that lasts as occurs with smoking, asthma or emphysema
■ cough that occurs with too much phlegm (mucus)

**Stop use and ask a doctor if cough lasts more than 7 days, cough comes
back, or occurs with fever, rash or headache that lasts. These could be
signs of a serious condition.**

**If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or
contact a Poison Control Center right away.**

Directions

■ **shake bottle well before use**

■ **dose as follows or as directed by a doctor**
adults and children 12 years of age and over 2 teaspoonfuls every 12 hours,
not to exceed 4 teaspoonfuls in 24 hours
children 6 to under 12 years of age 1 teaspoonful every 12 hours,
not to exceed 2 teaspoonfuls in 24 hours
children 2 to under 6 years of age 1/2 teaspoonful every 12 hours,
not to exceed 1 teaspoonful in 24 hours
children under 2 years of age consult a doctor

Other information

■ each 5 mL teaspoonful contains: sodium 6 mg
■ store at 20°-25°C (68°-77°F)

Inactive ingredients: citric acid anhydrous, D&C Red #33, edetate
disodium, ethylcellulose, FD&C Blue #1, Flavor, high fructose corn syrup,
methylparaben, polyethylene glycol 3350, polysorbate 80, propylene glycol,
propylparaben, purified water, sucrose, terepenth, vegetable oil, xanthan gum

Questions? 1-866-682-4639



021207
4000530

© 2007, Distributed by Adams Respiratory Therapeutics, Fort Worth, TX 76155

GLUE AREA
NO INK & COATING

12 Hour Cough Relief
Delsym

NDC 63824-172-63

12 Hour Cough Relief
Delsym
dextromethorphan polistirex
extended-release suspension
**COUGH
SUPPRESSANT**

12 Hour Cough Relief
Delsym
dextromethorphan polistirex
extended-release suspension
**COUGH
SUPPRESSANT**

12 Hour Cough Relief
Delsym
dextromethorphan polistirex
extended-release suspension
**COUGH
SUPPRESSANT**

**NEW GRAPE
FLAVOR**



**Contains
No Fever Reducer
or Pain Reliever**
Great Grape Taste

**Also Available
In Great
Orange Flavor**



Alcohol-free
89 mL (3 fl oz)

Grape-Flavored Liquid
for Children & Adults

Please visit our Web site:
www.delsym.com

**Contains
No Fever Reducer
or Pain Reliever**
Great Grape Taste

See back panel for full dosing directions.

DELSYM® DOSING SHAKE WELL BEFORE USE.	
Age (yr)	Dose (1 TSP = 5 mL)
Under 2	CONSULT DOCTOR
2 to under 6	1/2 TEASPOONFUL EVERY 12 HOURS
6 to under 12	1 TEASPOONFUL EVERY 12 HOURS

Lot No.:
Exp. Date:

3-63824-172-63-7

GLUE AREA

NO INK & COATING

Job Description:
Drawing #: #C03
Carton Size (Built)
Software: Adobe
Date Work Perfor
Colors Used: 4/C
Fonts Used: Frut
Hel
Additional Infor



FILE INFORMATION

Job Description: Delsym Pediatric Grape 3-oz. Label

Drawing #: Not Available

Label Size: 5-5/8"w x 2-1/8"h

Software: Adobe Illustrator 12

Date Work Performed: 02-13-07

Colors Used: PMS 527 + PMS 2738 + Process Yellow

Fonts Used: Frutiger, Helvetica Condensed,
Helvetica Neue Cond, Helvetica Neue Ext

Additional Information:

No Fever Reducer

Contains

Delsym
dextromethorphan polistirex
extended-release suspension
**COUGH
SUPPRESSANT**

DELSYM® DOSING
SHAKE WELL BEFORE USE.

Age (yr)

Dose
(1 TSP = 5 mL)

12 years
to adult

2 TEASPOONFULS
EVERY 12 HOURS

6 to
under 12

1 TEASPOONFUL
EVERY 12 HOURS

12 Hour Cough Relief



Delsym
dextromethorphan polistirex
extended-release suspension
**COUGH
SUPPRESSANT**

NDC 63824-171-75

TAMPER EVIDENT:
Do not use if the neckband
printed with is broken or missing.

12 Hour Cough Relief

12 Hour Cough Relief

Delsym 12 Hour Cough Relief

Drug Facts

Active ingredient (in each 5 mL teaspoonful) **Purpose**
Dextromethorphan polistirex equivalent to
30 mg dextromethorphan hydrobromide.....Cough suppressant

Use temporarily relieves cough due to minor throat and bronchial
irritation as may occur with the common cold or inhaled irritants

Warnings

Do not use if you are now taking a prescription monoamine oxidase
inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional
conditions, or Parkinson's disease), or for 2 weeks after stopping the

Drug Facts (continued)

MAOI drug. If you do not know if your prescription drug contains an
MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- chronic cough that lasts as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if cough lasts more than 7 days, cough
comes back, or occurs with fever, rash or headache that lasts. These
could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help
or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- **Shake bottle well before use.**

■ dose as follows or as directed by a doctor

adults and children

12 years of age

and over

2 teaspoonfuls every

12 hours, not to exceed

4 teaspoonfuls in 24 hours

children 6 to under

12 years of age

1 teaspoonful every

12 hours, not to exceed

2 teaspoonfuls in 24 hours

children 2 to under

6 years of age

1/2 teaspoonful every

12 hours, not to exceed

1 teaspoonful in 24 hours

children under

2 years of age

consult a doctor

Other information

- each 5 mL teaspoonful contains: sodium 6 mg

■ store at 20°-25° (68°-77°)

Drug Facts (continued)

Inactive ingredients

citric acid anhydrous,
D&C Red #33, edetate disodium, ethylcellulose,
FD&C Blue #1, flavor, high fructose corn syrup,
methylparaben, polyethylene glycol 3350, polysorbate 80,
propylene glycol, propylparaben, purified water, sucrose,
tragacanth, vegetable oil, xanthan gum

Questions? 1-866-682-4639

ADAMS
RESPIRATORY
THERAPEUTICS

021507
4000525

© 2007. Distributed by
Adams Respiratory Therapeutics, Fort Worth, TX 76155

LOT & EXP DATE AREA
NO INK



GLUE
AREA
NO INK
OR COAT

Delsym
dextromethorphan polistirex
extended-release suspension
**COUGH
SUPPRESSANT**

Alcohol-free
12 bottles
containing
15 mL each
PHYSICIAN SAMPLES —
NOT TO BE SOLD

GLUE
AREA
NO INK
OR COAT

2 1/2"

3 3/4"

Great Grape Taste

Contains
No Fever Reducer
or Pain Reliever

Delsym
dextromethorphan polistirex
extended-release suspension
**COUGH
SUPPRESSANT**

Delsym
DELSYM® DOSING
SHAKE WELL BEFORE USE.

Age (yr)
Under 2

Dose
(1 TSP = 5 mL)

Under 2
CONSULT
DOCTOR

**2 to
under 6**
EVERY 12 HOURS

**6 to
under 12**
EVERY 12 HOURS

See side panel for full dosing directions.

12 Hour Cough Relief



Delsym
dextromethorphan polistirex
extended-release suspension
**COUGH
SUPPRESSANT**

NDC 63824-172-75

TAMPER EVIDENT:
Do not use if the neckband
printed with is broken or missing.

12 Hour Cough Relief

Delsym 12 Hour Cough Relief

Drug Facts

Active ingredient (in each 5 mL teaspoonful) Purpose
Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide.....Cough suppressant

Use temporarily relieves cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI); certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the

Drug Facts (continued)

MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- chronic cough that lasts as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Inactive ingredients citric acid anhydrous, D&C Red #33, eddiate disodium, ethylcellulose, FD&C Blue #1, flavor, high fructose corn syrup, methylparaben, polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, saccharin, vegetable oil, xanthan gum.

Questions? 1-866-682-4639

Drug Facts (continued)

Directions

■ shake the bottle well before use	2 teaspoonfuls every 12 hours, not to exceed 4 teaspoonfuls in 24 hours
■ use as follows or as directed by a doctor	12 years of age and over
■ adults and children	children 6 to under 12 years of age
■ 12 years of age and over	children 2 to under 6 years of age
■ 12 years of age	children under 2 years of age
■ 12 years of age	consult a doctor

Other information

- each 5 mL (teaspoonful) contains: sodium 6 mg
- Store at 20°-25°C (68°-77°F)



© 2007, distributed by
Adams Respiratory Therapeutics, Fort Worth, TX 76165

021907
4000526

LOT & EXP DATE AREA
NO INK



Grape-Flavored
Liquid for
Children & Adults

Delsym
dextromethorphan polistirex
extended-release suspension
**COUGH
SUPPRESSANT**

Alcohol-free
12 bottles
containing
15 mL each
PHYSICIAN SAMPLES —
NOT TO BE SOLD

2 1/2"

3 3/4"

Cover

PHYSICIAN SAMPLE - NOT TO BE SOLD

Delsym
dextromethorphan polistirex
extended-release suspension

COUGH SUPPRESSANT

15 mL (1/2 fl oz)

TAMPER EVIDENT: Do not use if the neckband printed with is broken or missing.

Active ingredient
(in each 5 mL teaspoonful)
Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide.....suppressant

Purpose
User: temporarily relieves cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants.

Warnings: Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI). (Certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Back of Cover

Ask a doctor before use if you have chronic cough that lasts as occurs with smoking, asthma, or emphysema; cough that occurs with too much phlegm (mucus). Stop use and ask a doctor if cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
Shake bottle well before use. Dose as follows or as directed by a doctor.

adults and children 12 years of age and over	2 teaspoonfuls every 12 hours, not to exceed 4 teaspoonfuls in 24 hours
children 6 to under 12 years of age	1 teaspoonful every 12 hours, not to exceed 2 teaspoonfuls in 24 hours
children 2 to under 6 years of age	1/2 teaspoonful every 12 hours, not to exceed 1 teaspoonful in 24 hours
children under 2 years of age	consult a doctor

Base

PHYSICIAN SAMPLE - NOT TO BE SOLD

Delsym
dextromethorphan polistirex
extended-release suspension

COUGH SUPPRESSANT

15 mL (1/2 fl oz)

Other information: each 5 mL teaspoonful contains: sodium 6 mg. Store at 20°-25°C (68°-77°F).

Inactive ingredients: citric acid anhydrous, D & C Red #33, edetate disodium, ethylcellulose, FD & C Blue #1, flavor, high fructose corn syrup, methylparaben, polyethylene glycol 3350, polyacrylate 60, propylene glycol, propylparaben, purified water, sucrose, triacetin, vegetable oil, xanthan gum

Questions? 1-866-682-4639

© 2007. Distributed by Adams Respiratory Therapeutics Fort Worth, TX 76155

Lot No.:
Exp. Date:
022207 4000614

15 1/16"

4 1/16"

Hinge Area

Reseal Area

FILE INFORMATION

Job Description: Delsym Grape 15mL Expanded Content Label

Drawing #: Not Available

Label Size: 4-1/16"w x 15/16"h

Software: Adobe Illustrator 12