

Unwinding Direction

NDC 68180-398-01

Celecoxib Capsules

400 mg

PHARMACIST: PLEASE DISPENSE WITH MEDICATION GUIDE PROVIDED SEPARATELY

Rx only

LUPIN

100 Capsules

Each capsule contains celecoxib USP 400 mg.

Usual Dosage: See accompanying prescribing information.

Storage: Store at 25°C (77°F); excursions permitted to 15 to 30°C (59 to 86°F) [see USP Controlled Room Temperature].

Dispense in tight (USP), child resistant containers.

Manufactured for: **Lupin Pharmaceuticals, Inc.** Baltimore, Maryland 21202 United States

Manufactured by: **Lupin Limited** Goa 403 722 INDIA

Code No. GO/DRUGS/654 241029

Barcode: 681801398011

23 x 70 mm


Unvarnished Area

LOT NO. EXP:

Artwork Same Size
Size 180 x 85 mm (W x H)
Comp1- D:\Jobs\Lupin-Ltd\US\Submission\Label\Celecoxib Capsules\
Celecoxib-400mg-100Caps Feb-2015 .cdr
1st Laser Proof 26/02/2k15

Market/Customer :-	US	Location	Goa
Prepared On	26/02/2k15	Version No	1
Product name	Celecoxib-400mg-100Caps		
Material code	241029	Supersedes Material code	223236
Minimum Font Size	10	Barcode value	N368180398011
Dimensions	180 x 85 (W x H) mm	Barcode Type (Ex. NDC, PZN, EAN-13)	UPC(A)
Substrate	Chromo Art	Pharmacode value	NA
Varnish Type	—	Component	Label
Pantone Colours	 PANTONE 354 C  PANTONE 7527 C  PANTONE 375 C  Black  DELINE - DOES NOT PRINT  AREA FOR BATCH CODING		
Reason of Revision: Change in Text Matter			

Unwinding Direction



Artwork Same Size

Size 135 x 50 mm (W x H)

Comp1- D:\Jobs\Lupin-Ltd\US\Submission\Label\Celecoxib Capsules\

Celecoxib-400mg-60Caps Feb-2015 .cdr

1st Laser Proof 26/02/2k15

Market/Customer :-	US	Location	Goa
Prepared On	26/02/2k15	Version No	1
Product name	Celecoxib-400mg-60Caps		
Material code	241028	Supercedes Material code	223235
Minimum Font Size	7	Barcode value	N368180398073
Dimensions	135 x 50 (W x H) mm	Barcode Type (Ex. NDC, PZN, EAN-13)	UPC(A)
Substrate	Chromo Art	Pharmacode value	NA
Varnish Type	---	Component	Label
Pantone Colours	PANTONE 354 C PANTONE 7527 C	PANTONE 375 C Black	DIELINE - DOES NOT PRINT AREA FOR BATCH CODING
Reason of Revision: Change in Text Matter			

Unwinding Direction

NDC 68180-397-03

Celecoxib Capsules

200 mg

PHARMACIST: PLEASE DISPENSE WITH MEDICATION GUIDE PROVIDED SEPARATELY

Rx only

1000 Capsules

LUPIN

Each capsule contains celecoxib USP 200 mg.

Usual Dosage: See accompanying prescribing information.

Storage: Store at 25°C (77°F); excursions permitted to 15 to 30°C (59 to 86°F) [see USP Controlled Room Temperature].

Dispense in tight (USP), child resistant containers.

Manufactured for:

Lupin Pharmaceuticals, Inc.

Baltimore, Maryland 21202

United States

Manufactured by:

Lupin Limited

Goa 403 722 INDIA

Code No. GO/DRUGS/654 241027



32 x 74 mm

LOT NO. EXP:

Unvarnished Area

Artwork Same Size
 Size : 200 x 90 (L x W)
 Comp-1 D:\Jobs\Lupin Ltd\US\Submission\Label\Celecoxib Capsules\
 Celecoxib-200mg-1000Caps Feb-2015 .cdr
 1st Laser Proof 26/02/2k15

Market/Customer :-	US	Location	Goa
Prepared On	26/02/2k15	Version No	1
Product name	Celecoxib-200mg-1000Caps		
Material code	241027	Supersedes Material code	223234
Minimum Font Size	10	Barcode value	N368180397038
Dimensions	200 x 90 (W x H) mm	Barcode Type (Ex. NDC, PZN, EAN-13)	UPC(A)
Substrate	Chromo Art	Pharmacode value	NA
Varnish Type	—	Component	Label
Pantone Colours	PANTONE 1235 C PANTONE 7527 C PANTONE 375 C Black DIELINE - DOES NOT PRINT AREA FOR BATCH CODING		
Reason of Revision: Change in Text Matter			

DELINE - DOES NOT PRINT

AREA FOR BATCH CODING

Unwinding Direction



NDC 68180-397-02

Celecoxib Capsules

200 mg

PHARMACIST: PLEASE DISPENSE WITH MEDICATION GUIDE PROVIDED SEPARATELY

Rx only

LUPIN

Each capsule contains celecoxib USP 200 mg.

Usual Dosage: See accompanying prescribing information.

Storage: Store at 25°C (77°F); excursions permitted to 15 to 30°C (59 to 86°F) [see USP Controlled Room Temperature].

Dispense in tight (USP), child resistant containers.

Manufactured for:
Lupin Pharmaceuticals, Inc.
Baltimore, Maryland 21202
United States

Code No. GO/DRUGS/654

Manufactured by:
Lupin Limited
Goa 403 722 INDIA

241026

32 x 74 mm

LOT NO.
EXP:




68180-39702-1

Unvarnished Area

Artwork Same Size
Size : 200 x 90 (L x W)
Comp-1 D:\Jobs\Lupin Ltd\US\Submission\Label\Celecoxib Capsules\
Celecoxib-200mg-500Caps Feb-2015 .cdr
1st Laser Proof 26/02/2k15

Market/Customer :-	US	Location	Goa
Prepared On	26/02/2k15	Version No	1
Product name	Celecoxib-200mg-500Caps		
Material code	241026	Supersedes Material code	223233
Minimum Font Size	10	Barcode value	N368180397021
Dimensions	200 x 90 (W x H) mm	Barcode Type (Ex. NDC, PZN, EAN-13)	UPC(A)
Substrate	Chromo Art	Pharmacode value	NA
Varnish Type	—	Component	Label
Pantone Colours	 PANTONE 1235 C  PANTONE 7527 C  PANTONE 375 C  Black  DIE LINE - DOES NOT PRINT  AREA FOR BATCH CODING		
Reason of Revision: Change in Text Matter			

Unwinding Direction



Artwork Same Size
 Size 135 x 50 mm (W x H)
 Comp1- D:\Jobs\Lupin-Ltd\US\Submission\Label\Celecoxib Capsules\
 Celecoxib-200mg-100Caps Feb-2015 .cdr
 1st Laser Proof 26/02/2k15

Market/Customer :-	US	Location	Goa
Prepared On	26/02/2k15	Version No	1
Product name	Celecoxib-200mg-100Caps		
Material code	241025	Supersedes Material code	223232
Minimum Font Size	7	Barcode value	N368180397014
Dimensions	135 x 50 (W x H) mm	Barcode Type (Ex. NDC, PZN, EAN-13)	UPC(A)
Substrate	Chromo Art	Pharmacode value	NA
Varnish Type	—	Component	Label
Pantone Colours	PANTONE 1235 C PANTONE 7527 C	PANTONE 375 C Black	DIELINE - DOES NOT PRINT AREA FOR BATCH CODING
Reason of Revision: Change in Text Matter			

Unwinding Direction

The diagram shows a label for Celecoxib Capsules. A long arrow at the top points to the left, indicating the unwinding direction. The label itself is rectangular with a green vertical band on the left side. The text on the label includes: NDC 68180-396-03, Celecoxib Capsules, 100 mg, PHARMACIST: PLEASE DISPENSE WITH MEDICATION GUIDE PROVIDED SEPARATELY, Rx only, 1000 Capsules, LUPIN, Each capsule contains celecoxib USP 100 mg., Usual Dosage: See accompanying prescribing information., Storage: Store at 25°C (77°F); excursions permitted to 15 to 30°C (59 to 86°F) [see USP Controlled Room Temperature]. Dispense in tight (USP), child resistant containers., Manufactured for: Lupin Pharmaceuticals, Inc. Baltimore, Maryland 21202 United States, Manufactured by: Lupin Limited Goa 403 722 INDIA, Code No. GO/DRUGS/654 241024, and a barcode with the number 68180396031. A dashed box on the right side of the label indicates an 'Unvarnished Area' with dimensions 32 x 74 mm. The label also includes fields for LOT NO. and EXP.

Artwork Same Size
 Size : 200 x 90 (L x W)
 Comp-1 D:\Jobs\Lupin Ltd\US\Submission\Label\Celecoxib Capsules\
 Celecoxib-100mg-1000Caps Feb-2015 .cdr
 1st Laser Proof 26/02/2k15

Market/Customer :-	US	Location	Goa
Prepared On	26/02/2k15	Version No	1
Product name	Celecoxib-100mg-1000Caps		
Material code	241024	Supersedes Material code	223230
Minimum Font Size	10	Barcode value	N368180396031
Dimensions	200 x 90 (W x H) mm	Barcode Type (Ex. NDC, PZN, EAN-13)	UPC(A)
Substrate	Chromo Art	Pharmacode value	NA
Varnish Type	—	Component	Label
Pantone Colours	 PANTONE 2655 C  PANTONE 7527 C  PANTONE 375 C  Black  DIELINE - DOES NOT PRINT  AREA FOR BATCH CODING		
Reason of Revision: Change in Text Matter			

Unwinding Direction



NDC 68180-396-02

Celecoxib Capsules

100 mg

**PHARMACIST: PLEASE DISPENSE
WITH MEDICATION GUIDE
PROVIDED SEPARATELY**

Rx only

LUPIN

500 Capsules

Each capsule contains celecoxib
USP 100 mg.

Usual Dosage: See accompanying
prescribing information.

Storage: Store at 25°C (77°F); excursions
permitted to 15 to 30°C (59 to 86°F)
[see USP Controlled Room Temperature].
Dispense in tight (USP), child resistant
containers.

Manufactured for:
Lupin Pharmaceuticals, Inc.
Baltimore, Maryland 21202
United States

Manufactured by:
Lupin Limited
Goa 403 722 INDIA

Code No. GO/DRUGS/654

241023



N 68180396024

27 x 61.5 mm

LOT NO.


EXP.

Unvarnished Area

Artwork Same Size
Size 170 x 75 mm (W x H)
Comp1- D:\Jobs\Lupin-Ltd\US\Submission\Label\Celecoxib Capsules\
Celecoxib-100mg-500Caps Feb-2015 .cdr
1st Laser Proof 26/02/2k15

Market/Customer :-	US	Location	Goa
Prepared On	26/02/2k15	Version No	1
Product name	Celecoxib-100mg-500Caps		
Material code	241023	Supercedes Material code	223229
Minimum Font Size	9	Barcode value	N368180396024
Dimensions	170 x 75 (W x H) mm	Barcode Type (Ex, NDC, PZN, EAN-13)	UPC(A)
Substrate	Chromo Art	Pharmacode value	NA
Varnish Type	—	Component	Label
Pantone Colours	PANTONE 2655 C PANTONE 7527 C PANTONE 375 C Black DIELINE - DOES NOT PRINT AREA FOR BATCH CODING		
Reason of Revision: Change in Text Matter			


Unwinding Direction



Market/Customer >	US	Location	Goa
Prepared On	26/02/2k15	Version No	1
Product name	Celecoxib-100mg-100Caps		
Material code	241022	Supersedes Material code	223228
Minimum Font Size	4,5	Barcode value	N368180396017
Dimensions	90 x 37 (W x H) mm	Barcode Type (Ex. NDC, PZN, EAN-13)	UPC(A)
Substrate	Chromo Art	Pharmacode value	NA
Varnish Type	---	Component	Label
Pantone Colours	PANTONE 2855 C PANTONE 7527 C	PANTONE 375 C Black	DIELINE - DOES NOT PRINT AREA FOR BATCH CODING
Reason of Revision: Change in Text Matter			

Artwork Same Size
Size 90 x 37 mm (W x H)
Comp1- D:\Jobs\Lupin-Ltd\US\Submission\Label\Celecoxib Capsules\Celecoxib-100mg-100Caps Feb-2015 .cdr
1st Laser Proof 26/02/2k15

Unwinding Direction



Market/Customer :-	US	Location	Goa
Prepared On	28/02/2k15	Version No	1
Product name	Celecoxib-50mg-60Caps		
Material code	241021	Supersedes Material code	223196
Minimum Font Size	4.5	Barcode value	N366180395072
Dimensions	90 x 37 (W x H) mm	Barcode Type (Ex. NDC, PZN, EAN-13)	UPC(A)
Substrate	Chroma Art	Pharmacode value	NA
Varnish Type	---	Component	Label
Pantone Colours	PANTONE 220 C PANTONE 7527 C	PANTONE 375 C Black	DIELINE - DOES NOT PRINT AREA FOR BATCH CODING
Reason of Revision: Change in Text Matter			

Artwork Same Size
 Size 90 x 37 mm (W x H)
 Comp1- D:\Jobs\Lupin-Ltd\US\Submission\Label\Celecoxib Capsules\Celecoxib-50mg-60Caps Feb-2015 .cdr
 1st Laser Proof 26/02/2k15

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all of the information needed to use celecoxib capsules safely and effectively. See full prescribing information for celecoxib capsules.

CELECOXIB capsules, for oral use
Initial U.S. Approval: 1998

WARNING: CARDIOVASCULAR AND GASTROINTESTINAL RISKS	
See full prescribing information for complete boxed warning	
Cardiovascular Risk • Celecoxib capsules may cause an increased risk of serious cardiovascular thrombotic events, myocardial infarction, and stroke, which can be fatal. All NSAIDs may have a similar risk. This risk may increase with duration of use. Patients with cardiovascular disease or risk factors for cardiovascular disease may be at greater risk. (5.1,14.6) • Celecoxib capsules are contraindicated for the treatment of peri-operative pain in the setting of coronary artery bypass graft (CABG) surgery. (4, 5.1)	
Gastrointestinal Risk • NSAIDs, including celecoxib capsules, cause an increased risk of serious gastrointestinal adverse events including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients are at greater risk for serious gastrointestinal (GI) events. (5.4)	

INDICATIONS AND USAGE

- Celecoxib is a nonsteroidal anti-inflammatory drug indicated for:
- Osteoarthritis (OA) (1.1)
 - Rheumatoid Arthritis (RA) (1.2)
 - Juvenile Rheumatoid Arthritis (JRA) in patients 2 years and older (1.3)
 - Ankylosing Spondylitis (AS) (1.4)
 - Acute Pain (AP) (1.5)

DOSAGE AND ADMINISTRATION

- Use lowest effective dose for the shortest duration consistent with treatment goals for the individual patient. (1, 5.1, 5.4)
- OA: 200 mg once daily or 100 mg twice daily (2.1, 14.1)
 - RA: 100 to 200 mg twice daily (2.2, 14.2)
 - JRA: 50 mg twice daily in patients 10 to 25 kg, 100 mg twice daily in patients more than 25 kg (2.3, 14.3)
 - AS: 200 mg once daily single dose or 100 mg twice daily. If no effect is observed after 6 weeks, a trial of 400 mg (single or divided doses) may be of benefit (2.4, 14.4)
 - AP: 400 mg initially, followed by 200 mg dose if needed on first day. On subsequent days, 200 mg twice daily as needed (2.5, 14.5)
- Reduce daily dose by 50% in patients with moderate hepatic impairment (Child-Pugh Class B).
- Consider a dose reduction by 50% (or alternative management for JRA) in patients who are known or suspected to be CYP2C9 poor metabolizers, (2.6, 8.4, 8.8, 12.3).

FULL PRESCRIBING INFORMATION: CONTENTS*

BOXED WARNING

1 INDICATIONS AND USAGE

- 1.1 Osteoarthritis
- 1.2 Rheumatoid Arthritis
- 1.3 Juvenile Rheumatoid Arthritis
- 1.4 Ankylosing Spondylitis
- 1.5 Acute Pain

2 DOSAGE AND ADMINISTRATION

- 2.1 Osteoarthritis
- 2.2 Rheumatoid Arthritis
- 2.3 Juvenile Rheumatoid Arthritis
- 2.4 Ankylosing Spondylitis
- 2.5 Management of Acute Pain
- 2.6 Special Populations

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Cardiovascular Thrombotic Events
- 5.2 Hypertension
- 5.3 Congestive Heart Failure and Edema
- 5.4 Gastrointestinal (GI) Effects
- 5.5 Hepatic Effects
- 5.6 Renal Effects
- 5.7 Anaphylactoid Reactions
- 5.8 Skin Reactions
- 5.9 Pregnancy
- 5.10 Corticosteroid Treatment
- 5.11 Hematological Effects
- 5.12 Disseminated Intravascular Coagulation (DIC)
- 5.13 Preexisting Asthma
- 5.14 Laboratory Tests
- 5.15 Inflammation
- 5.16 Concomitant NSAID Use

6 ADVERSE REACTIONS

- 6.1 Pre-marketing Controlled Arthritis Trials
- 6.2 The Celecoxib Long-Term Arthritis Safety Study
- 6.3 Juvenile Rheumatoid Arthritis Study
- 6.4 Other Pre-Approval Studies
- 6.5 The APC and ProSAP Trials

7 DRUG INTERACTIONS

- 7.1 Warfarin
- 7.2 Lithium
- 7.3 Aspirin

FULL PRESCRIBING INFORMATION

WARNING: CARDIOVASCULAR AND GASTROINTESTINAL RISKS

- Cardiovascular Risk**
- Celecoxib capsules may cause an increased risk of serious cardiovascular thrombotic events, myocardial infarction, and stroke, which can be fatal. All nonsteroidal anti-inflammatory drugs (NSAIDs) may have a similar risk. This risk may increase with duration of use. Patients with cardiovascular disease or risk factors for cardiovascular disease may be at greater risk. (5.1,14.6)
 - Celecoxib capsules are contraindicated for the treatment of peri-operative pain in the setting of coronary artery bypass graft (CABG) surgery. (4, 5.1)
- Gastrointestinal Risk**
- NSAIDs, including celecoxib capsules, cause an increased risk of serious gastrointestinal adverse events including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients are at greater risk for serious gastrointestinal events. (5.4)

1 INDICATIONS AND USAGE

Carefully consider the potential benefits and risks of celecoxib capsules and other treatment options before deciding to use celecoxib capsules. Use the lowest effective dose for the shortest duration consistent with individual patient treatment goals (see WARNINGS AND PRECAUTIONS (5)).

1.1 Osteoarthritis (OA)

Celecoxib capsules are indicated for relief of the signs and symptoms of OA (see CLINICAL STUDIES (14.1)).

1.2 Rheumatoid Arthritis (RA)

Celecoxib capsules are indicated for relief of the signs and symptoms of RA (see CLINICAL STUDIES (14.2)).

1.3 Juvenile Rheumatoid Arthritis (JRA)

Celecoxib capsules are indicated for relief of the signs and symptoms of JRA in patients 2 years and older (see CLINICAL STUDIES (14.3)).

1.4 Ankylosing Spondylitis (AS)

Celecoxib capsules are indicated for the relief of signs and symptoms of AS (see CLINICAL STUDIES (14.4)).

1.5 Acute Pain (AP)

Celecoxib capsules are indicated for the management of AP in adults (see CLINICAL STUDIES (14.6)).

2. DOSAGE AND ADMINISTRATION

Use lowest effective dose for the shortest duration consistent with treatment goals for the individual patient. These doses can be given without regard to timing of meals.

2.1 Osteoarthritis

For relief of the signs and symptoms of OA the recommended oral dose is 200 mg per day administered as a single dose or as 100 mg twice daily.

2.2 Rheumatoid Arthritis

For relief of the signs and symptoms of RA the recommended oral dose is 100 to 200 mg twice daily.

2.3 Juvenile Rheumatoid Arthritis

For relief of the signs and symptoms of JRA the recommended oral dose for pediatric patients (age 2 years and older) is based on weight. For patients ≤10 kg to ≤25 kg the recommended dose is 50 mg twice daily. For patients >25 kg the recommended dose is 100 mg twice daily.

2.4 Ankylosing Spondylitis

For the management of the signs and symptoms of AS, the recommended dose of celecoxib capsules is 200 mg daily in single (once per day) or divided (twice per day) doses. If no effect is observed after 6 weeks, a trial of 400 mg daily may be worthwhile. If no effect is observed after 6 weeks on 400 mg daily, a response is not likely and consideration should be given to alternate treatment options.

2.5 Management of Acute Pain

The recommended dose of celecoxib capsules is 400 mg initially, followed by an additional 200 mg dose if needed on the first day. On subsequent days, the recommended dose is 200 mg twice daily as needed.

2.6 Special Populations

Hepatic Insufficiency

The daily recommended dose of celecoxib capsules in patients with moderate hepatic impairment (Child-Pugh Class B) should be reduced by 50%. The use of celecoxib capsules in patients with severe hepatic impairment is not recommended (see WARNINGS AND PRECAUTIONS (5.5), USE IN SPECIFIC POPULATIONS (8.4) and CLINICAL PHARMACOLOGY (12.3)).

Poor Metabolizers of CYP2C9 Substrates

Patients who are known or suspected to be poor CYP2C9 metabolizers based on genotype or previous history/experience with other CYP2C9 substrates (such as warfarin, phenytoin) should be administered celecoxib with caution. Consider starting treatment at half the lowest recommended dose in poor metabolizers (i.e. CYP2C9*3/*3). Consider using alternative management in JRA patients who are poor metabolizers (see USE IN SPECIFIC POPULATIONS (8.8), and CLINICAL PHARMACOLOGY (12.5)).

3. DOSAGE FORMS AND STRENGTHS

Capsules: 50 mg, 100 mg, 200 mg and 400 mg

4. CONTRAINDICATIONS

- Celecoxib is contraindicated in:
 - In patients with known hypersensitivity to celecoxib, aspirin, or other NSAIDs.
 - In patients who have demonstrated allergic-type reactions to sulfonamides.
 - In patients who have experienced asthma, urticaria, or allergic-type reactions after taking aspirin or other NSAIDs. Severe anaphylactoid reactions to NSAIDs, some of them fatal, have been reported in such patients (see WARNINGS AND PRECAUTIONS (5.7, 5.13)).
- For the treatment of peri-operative pain in the setting of coronary artery bypass graft (CABG) surgery (see WARNINGS AND PRECAUTIONS (5.1)).

5. WARNINGS AND PRECAUTIONS

5.1 Cardiovascular Thrombotic Events

Chronic use of celecoxib may cause an increased risk of serious adverse cardiovascular thrombotic events, myocardial infarction, and stroke, which can be fatal. In the APC (Adenoma Prevention with Celecoxib) trial, the hazard ratio for the composite endpoint of cardiovascular death, MI, or stroke was 3.4 (95% CI 1.4 to 8.5) for celecoxib 400 mg twice daily and 2.8 (95% CI 1.1 to 7.2) with celecoxib 200 mg twice daily compared to placebo. Cumulative rates for this composite endpoint over 3 years were 3.0% (20/671 subjects) and 2.5% (17/685 subjects), respectively, compared to 0.9% (6/679 subjects) with placebo treatment. The increases in both celecoxib dose groups versus placebo-treated patients were mainly due to an increased incidence of myocardial infarction (see CLINICAL STUDIES (14.6)).

All NSAIDs, both COX-2 selective and non-selective, may have a similar risk. Patients with known CV disease or risk factors for CV disease may be at greater risk. To minimize the potential risk for an adverse CV event in patients treated with celecoxib, the lowest effective dose should be used for the shortest duration consistent with individual patient treatment goals. Physicians and patients should remain alert for the development of such events, even in the absence of previous CV symptoms. Patients should be informed about the signs

and/or symptoms of serious CV toxicity and the steps to take if they occur.

There is no consistent evidence that concurrent use of aspirin mitigates the increased risk of serious CV thrombotic events associated with NSAID use. The concomitant use of aspirin and celecoxib does increase the risk of serious GI events (see WARNINGS AND PRECAUTIONS (5.4)).

Two large, controlled, clinical trials of a different COX-2 selective NSAID for the treatment of pain in the first 10 to 14 days following CABG surgery found an increased incidence of myocardial infarction and stroke (see CONTRAINDICATIONS (4)).

5.2 Hypertension

As with all NSAIDs, celecoxib can lead to the onset of new hypertension or worsening of preexisting hypertension, either of which may contribute to the increased incidence of CV events. Patients taking thiazides or loop diuretics may have impaired response to these therapies when taking NSAIDs. NSAIDs, including celecoxib, should be used with caution in patients with hypertension. Blood pressure should be monitored closely during the initiation of therapy with celecoxib and throughout the course of therapy. The rates of hypertension from the CLASS trial in the celecoxib, ibuprofen and diclofenac-treated patients were 2.4%, 4.2% and 2.9%, respectively (see CLINICAL STUDIES (14.6)).

5.3 Congestive Heart Failure and Edema

Fluid retention and edema have been observed in some patients taking NSAIDs, including celecoxib (see ADVERSE REACTIONS (6.1)). In the CLASS study (see CLINICAL STUDIES (14.6)), the Kaplan-Meier cumulative rates at 9 months of peripheral edema in patients on celecoxib 400 mg twice daily (4-fold and 2-fold the recommended OA and RA doses, respectively), ibuprofen 800 mg three times daily and diclofenac 75 mg twice daily were 4.5%, 6.9% and 4.7%, respectively. Celecoxib should be used with caution in patients with fluid retention or heart failure.

5.4 Gastrointestinal (GI) Effects

Risk of GI Ulceration, Bleeding, and Perforation

NSAIDs, including celecoxib, can cause serious gastrointestinal events including bleeding, ulceration, and perforation of the stomach, small intestine or large intestine, which can be fatal. These serious adverse events can occur at any time, with or without warning symptoms, in patients treated with NSAIDs. Only one in five patients who develop a serious upper GI adverse event on NSAID therapy is symptomatic. Complicated and symptomatic ulcer rates were 0.78% at nine months for all patients in the CLASS trial, and 2.19% for the subgroup on low-dose ASA. Patients 65 years of age and older had an incidence of 1.40% at nine months, 3.06% when also taking ASA (see CLINICAL STUDIES (14.6)). With longer duration of use of NSAIDs, there is a trend for increasing the likelihood of developing a serious GI event at some time during the course of therapy. However, even short-term therapy is not without risk.

NSAIDs should be prescribed with extreme caution in patients with a prior history of ulcer disease or gastrointestinal bleeding. Patients with a prior history of peptic ulcer disease and/or gastrointestinal bleeding who use NSAIDs have a greater than 10-fold increased risk for developing a GI bleed compared to patients with neither of these risk factors. Other factors that increase the risk of GI bleeding in patients treated with NSAIDs include concomitant use of oral corticosteroids or anticoagulants, longer duration of NSAID therapy, smoking, use of alcohol, older age, and poor general health status. Most spontaneous reports of fatal GI events are in elderly or debilitated patients and therefore special care should be taken in treating this population.

To minimize the potential risk for an adverse GI event, the lowest effective dose should be used for the shortest duration consistent with individual patient treatment goals. Physicians and patients should remain alert for signs and symptoms of GI ulceration and bleeding during celecoxib therapy and promptly initiate additional evaluation and treatment if a serious GI adverse event is suspected. For high-risk patients, alternate therapies that do not involve NSAIDs should be considered.

5.5 Hepatic Effects

NSAIDs, including celecoxib, may cause liver-associated enzymes may occur in up to 15% of patients taking NSAIDs, and notable elevations of ALT or AST (approximately 3 or more times the upper limit of normal) have been reported in approximately 1% of patients in clinical trials with NSAIDs. These laboratory abnormalities may progress, may remain unchanged, or may be transient with continuing therapy. Rare cases of severe hepatic reactions, including jaundice and fatal fulminant hepatitis, liver necrosis and hepatic failure (some with fatal outcome) have been reported with NSAIDs, including celecoxib (see ADVERSE REACTIONS (6.1)). In controlled clinical trials of celecoxib, the incidence of borderline elevations (greater than or equal to 1.2 times and less than 3 times the upper limit of normal) of liver associated enzymes was 6% for celecoxib and 5% for placebo, and approximately 0.2% of patients taking celecoxib and 0.3% of patients taking placebo had notable elevations of ALT and AST.

A patient with symptoms and/or signs suggesting liver dysfunction, or in whom an abnormal liver test has occurred, should be monitored carefully for evidence of the development of a more severe hepatic reaction while on therapy with celecoxib. If clinical signs and symptoms consistent with liver disease develop, or if systemic manifestations occur (e.g., eosinophilia, rash, etc.), celecoxib should be discontinued.

5.6 Renal Effects

Long-term administration of NSAIDs has resulted in renal papillary necrosis and other renal injury. Renal toxicity has also been seen in patients in whom renal prostaglandins have a compensatory role in the maintenance of renal perfusion. In these patients, administration of an NSAID may cause a dose-dependent reduction in prostaglandin formation and, secondarily, in renal blood flow, which may precipitate overt renal decompensation. Patients at greatest risk of this reaction are those with impaired renal function, heart failure, liver dysfunction, those taking diuretics, ACE-inhibitors, angiotensin II receptor antagonists, and the elderly. Discontinuation of NSAID therapy is usually followed by recovery to the pretreatment state. Clinical trials with celecoxib have shown renal effects similar to those observed with comparator NSAIDs.

No information is available from controlled clinical studies regarding the use of celecoxib in patients with advanced renal disease. Therefore, treatment with celecoxib is not recommended in these patients with advanced renal disease. If celecoxib therapy must be initiated, close monitoring of the patient's renal function is advisable.

5.7 Anaphylactoid Reactions

As with NSAIDs in general, anaphylactoid reactions have occurred in patients without known prior exposure to celecoxib. In post-marketing experience, rare cases of anaphylactic reactions and angioedema have been reported in patients receiving celecoxib. Celecoxib should not be given to patients with the aspirin triad. This symptom complex typically occurs in asthmatic patients who experience rhinitis with or without nasal polyps, or who exhibit severe, potentially fatal bronchospasm after taking aspirin or other NSAIDs (see CONTRAINDICATIONS (4), WARNINGS AND PRECAUTIONS (5.7)). Emergency help should be sought in cases where an anaphylactoid reaction occurs.

5.8 Skin Reactions

Celecoxib is a sulfonamide and can cause serious skin adverse events such as exfoliative dermatitis, Stevens-Johnson syndrome (SJS), and toxic epidermal necrolysis (TEN), which can be fatal. These serious events can occur without warning and in patients without prior known sufa allergy. Patients should be informed about the signs and symptoms of serious skin manifestations and the use of the drug should be discontinued at the first appearance of skin rash or any other sign of hypersensitivity.

5.9 Pregnancy

In late pregnancy, starting at 30 weeks gestation, celecoxib should be avoided because it may cause premature closure of the ductus arteriosus (see USE IN SPECIFIC POPULATIONS (8.1)).

5.10 Corticosteroid Treatment

Celecoxib cannot be expected to substitute for corticosteroids or to treat corticosteroid insufficiency. Abrupt discontinuation of corticosteroids may lead to exacerbation of corticosteroid-responsive illnesses. Patients on prolonged corticosteroid therapy should have their therapy tapered slowly if a decision is made to discontinue corticosteroids.

Capsules: 50 mg, 100 mg, 200 mg and 400 mg (3)

DOSAGE FORMS AND STRENGTHS

CONTRAINDICATIONS

- Known hypersensitivity to celecoxib or sulfonamides (4)
- History of asthma, urticaria, or other allergic-type reactions after taking aspirin or other NSAIDs (4, 5.7, 5.8, 5.13)
- Use during the perioperative period in the setting of coronary artery bypass graft (CABG) surgery (4, 5.1)

WARNINGS AND PRECAUTIONS

- Serious and potentially fatal cardiovascular (CV) thrombotic events, myocardial infarction, and stroke. Patients with known CV disease/risk factors may be at greater risk (5.1, 14.6, 17.2).
- Serious gastrointestinal (GI) adverse events, which can be fatal. The risk is greater in patients with a prior history of ulcer disease or GI bleeding, and in patients at high risk for GI events, especially the elderly. Celecoxib should be used with caution in these patients (5.4, 8.5, 14.6, 17.3).
- Elevated liver enzymes and, rarely, severe hepatic reactions. Discontinue use of celecoxib immediately if abnormal liver enzymes persist or worsen (5.5, 17.4).
- New onset or worsening of hypertension. Blood pressure should be monitored closely during treatment with celecoxib (5.2, 7.4, 17.2).
- Fluid retention and edema. Celecoxib should be used with caution in patients with fluid retention or heart failure (5.3, 17.6).
- Renal papillary necrosis and other renal injury with long term use. Use celecoxib with caution in the elderly, those with impaired renal function, heart failure, liver dysfunction, and those taking diuretics, ACE-inhibitors, or angiotensin II antagonists (5.6, 7.4, 8.7, 17.6).
- Anaphylactoid reactions. Do not use celecoxib in patients with the aspirin triad (5.7, 10, 17.7).
- Serious skin adverse events such as exfoliative dermatitis, Stevens-Johnson syndrome (SJS), and toxic epidermal necrolysis (TEN), which can be fatal and can occur without warning even without known prior sufa allergy. Discontinue celecoxib at first appearance of rash or skin reactions (5.8, 17.5).

ADVERSE REACTIONS

Most common adverse reactions in arthritis trials (>2% and >placebo): abdominal pain, diarrhea, dyspepsia, flatulence, peripheral edema, accidental injury, dizziness, pharyngitis, rhinitis, sinusitis, upper respiratory tract infection, rash (6.1).

To report SUSPECTED ADVERSE REACTIONS, contact Lupin Pharmaceuticals, Inc at 1-800-309-2561 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

DRUG INTERACTIONS

- Concomitant use of celecoxib and warfarin may result in increased risk of bleeding complications. (7.1)
- Concomitant use of celecoxib increases lithium plasma levels. (7.2)
- Concomitant use of celecoxib may reduce the antihypertensive effect of ACE Inhibitors and angiotensin II antagonists (7.4)
- Use caution with drugs known to inhibit P450 2C9 or metabolized by 2D6 due to the potential for increased plasma levels (2.6, 8.4, 8.8, 12.3)

USE IN SPECIFIC POPULATIONS

- Pregnancy Category C prior to 30 weeks gestation; Category D starting at 30 weeks gestation (5.9, 8.1, 17.8)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide

- 7.4 ACE-Inhibitors and Angiotensin II Antagonists
- 7.5 Fluconazole
- 7.6 Furosemide
- 7.7 Methotrexate
- 7.8 Concomitant NSAID Use

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Labor and Delivery
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Hepatic Insufficiency
- 8.7 Renal Insufficiency
- 8.8 Poor Metabolizers of CYP2C9 Substrates

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics
- 12.5 Pharmacogenomics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 13.2 Animal Toxicology

14 CLINICAL STUDIES

- 14.1 Osteoarthritis
- 14.2 Rheumatoid Arthritis
- 14.3 Juvenile Rheumatoid Arthritis
- 14.4 Ankylosing Spondylitis
- 14.5 Analgesia
- 14.6 Special Studies

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

- 17.1 Medication Guide
- 17.2 Cardiovascular Effects
- 17.3 Gastrointestinal Effects
- 17.4 Hepatic Effects
- 17.5 Adverse Skin Reactions
- 17.6 Weight Gain and Edema
- 17.7 Anaphylactoid Reactions
- 17.8 Effects During Pregnancy
- 17.9 Preexisting Asthma

* Sections or subsections omitted from the Full Prescribing Information are not listed.

5.11 Hematological Effects

Anemia is sometimes seen in patients receiving celecoxib. In controlled clinical trials the incidence of anemia was 0.6% with celecoxib and 0.4% with placebo. Patients on long-term treatment with celecoxib should have their hemoglobin or hematocrit checked if they exhibit any signs or symptoms of anemia or blood loss. Celecoxib does not generally affect platelet counts, prothrombin time (PT), or partial thromboplastin time (PTT), and does not inhibit platelet aggregation at indicated dosages (see CLINICAL PHARMACOLOGY (12.3)).

5.12 Disseminated Intravascular Coagulation (DIC)

Celecoxib should be used only with caution in pediatric patients with systemic onset JRA due to the risk of disseminated intravascular coagulation.

5.13 Preexisting Asthma

Patients with asthma may have aspirin-sensitive asthma. The use of aspirin in patients with aspirin-sensitive asthma has been associated with severe bronchospasm, which can be fatal. Since cross reactivity, including bronchospasm, between aspirin and other nonsteroidal anti-inflammatory drugs has been reported in such aspirin-sensitive patients, celecoxib should not be administered to patients with this form of aspirin sensitivity and should be used with caution in patients with preexisting asthma.

5.14 Laboratory Tests

Because serious GI tract ulcerations and bleeding can occur without warning symptoms, physicians should monitor for signs or symptoms of GI bleeding. Patients on long-term treatment with NSAIDs should have a CBC and a chemistry profile checked periodically. If abnormal liver tests or renal tests persist or worsen, celecoxib should be discontinued.

In controlled clinical trials, elevated BUN occurred more frequently in patients receiving celecoxib compared with patients on placebo. This laboratory abnormality was also seen in patients who received comparator NSAIDs in these studies. The clinical significance of this abnormality has not been established.

5.15 Inflammation

The pharmacological activity of celecoxib in reducing inflammation, and possibly fever, may diminish the utility of these diagnostic signs in detecting infectious complications of presumed noninfectious, painful conditions.

5.16 Concomitant NSAID Use

The concomitant use of celecoxib with any dose of a non-aspirin NSAID should be avoided due to the potential for increased risk of adverse reactions.

6. ADVERSE REACTIONS

Of the celecoxib-treated patients in the pre-marketing controlled clinical trials, approximately 4,250 were patients with OA, approximately 2,100 were patients with RA, and approximately 1,050 were patients with post-surgical pain. More than 8,500 patients received a total daily dose of celecoxib of 200 mg (100 mg twice daily or 200 mg once daily) or more, including more than 400 treated at 800 mg (400 mg twice daily). Approximately 3,900 patients received celecoxib at these doses for 6 months or more; approximately 2,300 of these have received it for 1 year or more and 124 of these have received it for 2 years or more. Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice. The adverse reaction information from clinical trials does, however, provide a basis for identifying the adverse events that appear to be related to drug use and for approximating rates.

6.1 Pre-marketing Controlled Arthritis Trials

Table 1 lists all adverse events, regardless of causality, occurring in ≥ 2% of patients receiving celecoxib from 12 controlled studies conducted in patients with OA or RA that included a placebo and/or a positive control group. Since these 12 trials were of different durations, and patients in the trials may not have been exposed for the same duration of time, these percentages do not capture cumulative rates of occurrence.

Table 1: Adverse Events Occurring in ≥ 2% of Celecoxib Patients from Pre-marketing Controlled Arthritis Trials					
	CeX N=4146	Placebo N=1854	NAP N=1366	DCF N=387	IBU N=345
Gastrointestinal					
Abdominal Pain	4.1%	2.8%	7.7%	9.0%	9.0%
Diarrhea	5.6%	3.6%	5.3%	9.3%	5.8%
Dyspepsia	8.8%	6.2%	12.2%	10.9%	12.8%
Flatulence	2.2%	1.0%	3.6%	4.1%	3.5%
Nausea	3.5%	4.2%	6.0%	3.4%	6.7%
Body as a whole					
Back Pain	2.8%	3.6%	2.2%	2.6%	0.9%
Peripheral Edema	2.1%	1.1%	2.1%	1.0%	3.5%
Injury, Accidental	2.9%	2.3%	3.0%	2.6%	3.2%
Central, Peripheral Nervous system					
Dizziness	2.0%	1.7%	2.6%	1.3%	2.3%
Headache	15.8%	20.2%	14.5%	15.5%	15.4%
Psychiatric					
Insomnia	2.3%	2.3%	2.9%	1.3%	1.4%
Respiratory					
Pharyngitis	2.3%	1.1%	1.7%	1.6%	2.6%
Rhinitis	2.0%	1.3%	2.4%	2.3%	0.6%
Sinusitis	5.0%	4.9%	4.0%	5.4%	5.8%
Upper Respiratory Infection	8.1%	6.7%	9.9%	9.8%	9.9%
Skin					
Rash	2.2%	2.1%	2.1%	1.3%	1.2%

Table 2: Adverse Events Occurring in ≥5% of JRA Patients in Any Treatment Group, by System Organ Class (of patients with events)

System Organ Class Preferred Term	All Doses Twice Daily		
	Celecoxib 3 mg/kg N=77	Celecoxib 6 mg/kg N=82	Naproxen 7.5 mg/kg N=83
Any Event	64	70	72
Eye Disorders	5	5	5
Gastrointestinal	26	24	36
Abdominal pain NOS	4	7	7
Abdominal pain upper	8	6	11
Vomiting NOS	1	1	1
Diarrhea NOS	5	4	8
Nausea	7	4	11
General	13	11	18
Pyrrexia	8	9	11
Infections	26	20	27
Nasopharyngitis	5	6	5
Injury and Poisoning	4	6	5
Investigations*	3	11	7
Musculoskeletal	8	10	17
Arthralgia	3	7	4
Nervous System	17	11	21
Headache NOS	13	10	16
Dizziness (extl vertigo)	1	1	7
Respiratory	8	15	16
Cough	7	7	8
Skin & Subcutaneous	10	7	18

* Abnormal laboratory tests, which include: Prolonged activated partial thromboplastin time, Bacteriuria NOS present, Blood creatine phosphokinase increased, Blood culture positive, Blood glucose increased, Blood pressure increased, Blood uric acid increased, Hematocrit decreased, Hematuna present, Hemoglobin decreased, Liver function tests NOS abnormal, Protenuria present, Transaminase NOS increased, Urine analysis abnormal NOS

6.4 Other Pre-Approval Studies

Adverse Events from Analyzing Spondylitis Studies
A total of 378 patients were treated with celecoxib in placebo- and active-controlled AS studies. Doses up to 400 mg once daily were studied. The types of adverse events reported in the AS studies were similar to those reported in the OA/RA studies.

Adverse Events from Analgesia Studies

Approximately 1,700 patients were treated with celecoxib in analgesia studies. All patients in post-oral surgery pain studies received a single dose of study medication. Doses up to 600 mg/day of celecoxib were studied in post-orthopedic surgery pain studies. The types of adverse events in the analgesia studies were similar to those reported in arthritis studies. The only additional adverse event reported was post-dental extraction alveolar osteitis (dry socket) in the post-oral surgery pain studies.

6.5 The APC and PreSAP Trials

Adverse Reactions from Long-term, Placebo-Controlled PVP Prevention Studies
Exposure to celecoxib in the APC and PreSAP trials was due to 800 mg daily for up to 3 years. [See *Special Studies Adenomatous Polyp Prevention Studies* (14.6)].

Some adverse reactions occurred in higher percentages of patients than in the arthritis pre-marketing trials (treatment durations up to 12 weeks; see *Adverse events from celecoxib pre-marketing controlled arthritis trials*, above). The adverse reactions for which these differences in patients treated with celecoxib were greater as compared to the arthritis pre-marketing trials were as follows:

	Celecoxib Capsules (400 to 800 mg daily) N = 2285	Placebo N=1303
Diarrhea	10.5%	7.0%
Gastroesophageal reflux disease	4.7%	3.1%
Nausea	6.8%	5.3%
Vomiting	3.2%	2.1%
Dyspnea	2.8%	1.6%
Hypertension	12.5%	9.9%

The following additional adverse reactions occurred in 20.1% and <1% of patients taking celecoxib capsules, at an incidence greater than placebo in the long-term polyp prevention studies, and were either not reported during the controlled arthritis pre-marketing trials or occurred with greater frequency in the long-term, placebo-controlled polyp prevention studies:

Adverse system disorders: Cervical infection
Eye disorders: Vitreous floaters, conjunctival hemorrhage
Fat and labyrinth: Labyrinthitis

Cardiac disorders: Angina unstable, aortic valve incompetence, coronary artery atherosclerosis, sinus bradycardia, ventricular hypertrophy
Vascular disorders: Deep vein thrombosis

Reproductive system and breast disorders: Ovarian cyst
Investigations: Blood potassium increased, blood sodium increased, blood testosterone decreased
Injury, poisoning and procedural complications: Epicondylitis, tendon rupture

7. DRUG INTERACTIONS

General
Celecoxib metabolism is predominantly mediated via cytochrome P450 (CYP) 2C9 in the liver. Co-administration of celecoxib with drugs that are known to inhibit CYP2C9 should be used with caution. Significant interactions may occur when celecoxib is administered together with drugs that inhibit CYP2C9. *In vitro* studies indicate that celecoxib, although not a substrate, is an inhibitor of CYP2C6. Therefore, there is a potential for an *in vivo* drug interaction with drugs that are metabolized by CYP2C6.

7.1 Warfarin

Anticoagulant activity should be monitored, particularly in the first few days, after initiating or changing celecoxib therapy in patients receiving warfarin or similar agents, since these patients are at an increased risk of bleeding complications. The effect of celecoxib on the anticoagulant effect of warfarin was studied in a group of healthy subjects receiving daily 2 to 5 mg doses of warfarin. In these subjects, celecoxib did not alter the anticoagulant effect of warfarin as determined by prothrombin time. However, in post-marketing experience, serious bleeding events, some of which were fatal, have been reported, predominantly in the elderly, in association with increases in prothrombin time in patients receiving celecoxib concurrently with warfarin.

7.2 Lithium

In a study conducted in healthy subjects, mean steady-state lithium plasma levels increased approximately 17% in subjects receiving lithium 450 mg twice daily with celecoxib 200 mg twice daily as compared to subjects receiving lithium alone. Patients on lithium treatment should be closely monitored when celecoxib is introduced or withdrawn.

7.3 Aspirin

Celecoxib can be used with low-dose aspirin. However, concomitant administration of aspirin with celecoxib increases the rate of GI ulceration or other complications, compared to use of celecoxib alone [see *WARN INGS AND PRECAUTIONS* (5.1, 5.4) and *CLINICAL STUDIES* (14.6)]. Because of its lack of platelet effects, celecoxib is not a substitute for aspirin for cardiovascular prophylaxis [see *CLINICAL PHARMACOLOGY* (12.2)].

7.4 ACE-inhibitors and Angiotensin II Antagonists

Reports suggest that NSAIDs may diminish the antihypertensive effect of Angiotensin Converting Enzyme (ACE) inhibitors and angiotensin II antagonists. This interaction should be given consideration in patients taking celecoxib concurrently with ACE-inhibitors and angiotensin II antagonists [see *CLINICAL PHARMACOLOGY* (12.2)].

7.5 Fluconazole

Concomitant administration of fluconazole at 200 mg once daily resulted in a two-fold increase in celecoxib plasma concentration. The inhibition of celecoxib metabolism by fluconazole was P450 2C9 by fluconazole [see *CLINICAL PHARMACOLOGY* (12.3)]. Celecoxib should be introduced at the lowest recommended dose in patients receiving fluconazole.

7.6 Furosemide

Clinical studies, as well as post-marketing observations, have shown that NSAIDs can reduce the natriuretic effect of furosemide and thiazides in some patients. This response has been attributed to inhibition of renal prostaglandin synthesis.

7.7 Methotrexate

In an interaction study of rheumatoid arthritis patients taking methotrexate, celecoxib did not have an effect on pharmacokinetics of methotrexate [see *CLINICAL PHARMACOLOGY* (12.3)].

7.8 Concomitant NSAID Use

The concomitant use of celecoxib with any dose of a non-aspirin NSAID should be avoided due to the potential for increased risk of adverse reactions.

8. USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category C: Pregnancy category D from 30 weeks of gestation onward.

Teratogenic Effects

Celecoxib at oral doses ≥150 mg/kg/day (approximately 2-fold human exposure at 200 mg twice daily as measured by AUC₀₋₁₂) caused an increased incidence of ventricular septal defects, a rare event, and fetal alterations, such as ribs fused, sternbrae fused and sternbrae misshapen when rabbits were treated throughout organogenesis. A dose-dependent increase in diaphragmatic hernias was observed when rats were given celecoxib at oral doses ≥30 mg/kg/day (approximately 6-fold human exposure based on the AUC₀₋₁₂ at 200 mg twice daily) throughout organogenesis. There are no studies in pregnant women. Celecoxib should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nonteratogenic Effects

Celecoxib produced pre-implantation and post-implantation losses and reduced embryo/fetal survival in rats at oral doses ≥50 mg/kg/day (approximately 6-fold human exposure based on the AUC₀₋₁₂ at 200 mg twice daily). These changes are expected with inhibition of prostaglandin synthesis and are not the result of permanent alteration of female reproductive function, nor are they expected at clinical exposures. No studies have been conducted to evaluate the effect of celecoxib on the closure of the ductus arteriosus in humans. Therefore, use of celecoxib during the third trimester of pregnancy should be avoided.

8.2 Labor and Delivery

Celecoxib produced no evidence of delayed labor or parturition at oral doses up to 100 mg/kg in rats (approximately 7-fold human exposure as measured by the AUC₀₋₁₂ at 200 mg BID). The effects of celecoxib on labor and delivery in pregnant women are unknown.

8.3 Nursing Mothers

From 3 published reports that included a total of 12 breastfeeding women showed low levels of celecoxib in breast milk. The calculated average daily infant dose was 10 to 40 mg/kg/day, less than 1% of the weight-based therapeutic dose for a two-year-old child. A report of two breastfed infants 17 and 22 months of age did not show any adverse events. Caution should be exercised when celecoxib is administered to a nursing woman.

8.4 Pediatric Use

Celecoxib is approved for relief of the signs and symptoms of Juvenile Rheumatoid Arthritis in patients 2 years and older. Safety and efficacy have not been studied beyond six months in children. The long-term cardiovascular toxicity in children exposed to celecoxib has not been evaluated and it is unknown if long-term risks may be similar to that seen in adults exposed to celecoxib or other COX-2 selective and non-selective NSAIDs [see *BOXED WARNING, WARN INGS AND PRECAUTIONS* (5.12), and *CLINICAL STUDIES* (14.3)].

The use of celecoxib in patients 2 years to 17 years of age with psoriatic/arthritis, polyarthralgia course JRA or in patients with systemic onset JRA was studied in a 12-week, double-blind, active controlled, pharmacokinetic, safety and efficacy study, with a 12-week open-label extension. Celecoxib has not been studied in children under the age of 2 years, in patients with body weight less than 10 kg (22 lbs), and in patients with active systemic features. Patients with systemic onset JRA (without active systemic features) appear to be at risk for the development of abnormal coagulation laboratory tests. In some patients with systemic onset JRA, celecoxib and naproxen were associated with mild prolongation of activated partial thromboplastin time (APTT) but not prothrombin time (PT). NSAIDs including celecoxib should be used only with caution in patients with systemic onset JRA, due to the risk of disseminated intravascular coagulation. Patients with systemic onset JRA should be monitored for the development of abnormal coagulation tests [see *DOSEAGE AND ADMINISTRATION* (2.3), *WARNINGS AND PRECAUTIONS* (5.12), *ADVERSE REACTIONS* (5.3), *ANIMAL TOXICOLOGY STUDIES* (14.3), and *CLINICAL STUDIES* (14.3)].

Alternative therapies for treatment of JRA should be considered in pediatric patients identified to be CYP2C9 poor metabolizers [see *Poor Metabolizers of CYP2C9 Substrates* (8.3)].

8.5 Geriatric Use

Of the total number of patients who received celecoxib in pre-approval clinical trials, more than 3,900 were 65 to 74 years of age, while approximately 1,300 additional patients were 75 years and over. No substantial differences in effectiveness were observed between these subjects and younger subjects. In clinical studies comparing renal function as measured by the GFR, BUN and creatinine, and platelet function as measured by bleeding time and platelet aggregation, the results were not different between elderly and young volunteers. However, as with other NSAIDs, elderly subjects that selectively inhibit COX-2, there have been more spontaneous post-marketing reports of fatal GI events and acute renal failure in the elderly than in younger patients [see *WARN INGS AND PRECAUTIONS* (5.4, 5.6)].

8.6 Hepatic Insufficiency

The daily recommended dose of celecoxib capsules in patients with moderate hepatic impairment (Child-Pugh Class B) should be reduced by 50%. The use of celecoxib in patients with severe hepatic impairment is not recommended [see *DOSEAGE AND ADMINISTRATION* (2.6) and *CLINICAL PHARMACOLOGY* (12.3)].

8.7 Renal Insufficiency

Celecoxib is not recommended in patients with severe renal insufficiency [see *WARNINGS AND PRECAUTIONS* (5.6) and *CLINICAL PHARMACOLOGY* (12.3)].

8.8 Poor Metabolizers of CYP2C9 Substrates

Patients who are known or suspected to be poor CYP2C9 metabolizers based on genotype or previous history/experience with other CYP2C9 substrates (such as warfarin, phenytoin) should be administered celecoxib with caution. Consider starting treatment at half the lowest recommended dose in poor metabolizers (i.e., CYP2C9*3/3). Alternative management should be considered in JRA patients identified to be CYP2C9 poor metabolizers [see *DOSEAGE AND ADMINISTRATION* (2.6) and *CLINICAL PHARMACOLOGY* (12.3)].


10. OVERDOSEAGE

No overdose of celecoxib were reported during clinical trials. Doses up to 2400 mg/day for up to 10 days in 12 patients did not result in serious toxicity. Symptoms following acute NSAID overdoses are usually limited to lethargy, drowsiness, nausea, vomiting, and epigastric pain, which are generally reversible with supportive care. Gastrointestinal bleeding can occur. Hypertension, acute renal failure, respiratory depression and coma may occur, but are rare. Anaphylactoid reactions have been reported with therapeutic ingestion of NSAIDs, and may occur following an overdose.

Patients should be managed by symptomatic and supportive care following an NSAID overdose. There are no specific antidotes. No information is available regarding the removal of celecoxib by hemodialysis, but based on its high degree of plasma protein binding (>97%) dialysis is unlikely to be useful in overdose. Emesis and/or activated charcoal (60 to 100 g in adults, 1 to 2 g/kg in children) and/or osmotic cathartic may be indicated in patients seen within 4 hours of ingestion with symptoms or following a large overdose. Forced diuresis, alkalization of urine, hemodialysis, or hemoperfusion may not be useful due to high protein binding.

11. DESCRIPTION

Celecoxib is chemically designated as 4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl] benzenesulfonamide and is a diaryl-substituted pyrazole. The empirical formula is C₁₈H₁₅N₃O₃S, and the molecular weight is 381.38, the chemical structure is as follows:



Celecoxib oral capsules contain either 50 mg, 100 mg, 200 mg or 400 mg of celecoxib, together with inactive ingredients including black iron oxide, croscarmellose sodium, FD&C blue #1, FD&C red #40, gelatin, lactose monohydrate, magnesium stearate, potassium hydroxide, povidone, propylene glycol, shellac, sodium lauryl sulfate, titanium dioxide, red iron oxide and yellow iron oxide.

12. CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Celecoxib is a nonsteroidal anti-inflammatory drug that exhibits anti-inflammatory, analgesic, and antipyretic activities in animal models. The mechanism of action of celecoxib is believed to be due to inhibition of prostaglandin synthesis, primarily via inhibition of cyclooxygenase-2 (COX-2), and at therapeutic concentrations in humans, celecoxib does not inhibit the cyclooxygenase-1 (COX-1) isoenzyme. In animal colon tumor models, celecoxib reduced the incidence and multiplicity of tumors.

12.2 Pharmacodynamics

Platelets

In clinical trials using normal volunteers, celecoxib at single doses up to 800 mg and multiple doses of 600 mg twice daily for up to 7 days duration (higher than recommended therapeutic doses) had no effect on reduction of platelet aggregation or increase in bleeding time. Because of its lack of platelet effects, celecoxib is not a substitute for aspirin for cardiovascular prophylaxis. It is not known if there are any effects of celecoxib on platelets that may contribute to the increased risk of serious cardiovascular thrombotic adverse events associated with the use of celecoxib.

Fluid Retention

Inhibition of PGE2 synthesis may lead to sodium and water retention through increased reabsorption in the renal medullary thick ascending loop of Henle and perhaps other segments of the distal nephron. In the collecting ducts, PGE2 appears to inhibit water reabsorption by contracting the action of aldosterone hormone.

12.3 Pharmacokinetics

Absorption

Peak plasma levels of celecoxib occur approximately 3 hrs after an oral dose. Under fasting conditions, both peak plasma levels (C_{max}) and area under the curve (AUC) are roughly dose-proportional up to 200 mg B.D. at higher doses there are less than proportional increases in C_{max} and AUC [see *Food Effects*]. Absolute bioavailability studies have not been conducted. With multiple dosing, steady-state conditions are reached on or before Day 5. The pharmacokinetic parameters of celecoxib in a group of healthy subjects are shown in Table 3.

Table 3 Summary of Single Dose (200 mg) Disposition Kinetics of Celecoxib in Healthy Subjects*					
Mean (±SD) PK Parameter Values					
C _{max} , mg/mL	T _{max} , hr	Effective t _{1/2} , hr	V _d /F, L	CL/F, L/hr	
705 (38)	2.8 (37)	11.2 (31)	429 (34)	27.7 (28)	

*Subjects under fasting conditions (n=36, 19 to 52 yrs.)

Food Effects

When celecoxib capsules were taken with a high fat meal, peak plasma levels were delayed for about 1 to 2 hours with an increase in total absorption (AUC) of 10% to 20%. Under fasting conditions of platesd above 200 mg, there is less than a proportional increase in C_{max} and AUC, which is thought to be due to the low solubility of the drug in aqueous media.

Co-administration of celecoxib with an aluminum- and magnesium-containing antacid resulted in a reduction in plasma celecoxib concentrations with a decrease of 37% in C_{max} and 10% in AUC. Celecoxib at doses up to 200 mg twice daily, can be administered without regard to timing of meals. Higher doses (400 mg twice daily) should be administered with food to improve absorption.

In healthy adult volunteers, the overall systemic exposure (AUC) of celecoxib was equivalent when celecoxib was administered as intact capsule or capsule contents sprinkled on applesauce. There were no significant alterations in C_{max}, T_{max} or t_{1/2} after administration of capsule contents on applesauce [see *DOSEAGE AND ADMINISTRATION* (2)].

Distribution

In healthy subjects, celecoxib is highly protein bound (~97%) within the clinical dose range. *In vitro* studies indicate that celecoxib binds primarily to albumin and, to a lesser extent, α₂-acid glycoprotein. The apparent volume of distribution at steady state (V_d/F) is approximately 400 L, suggesting extensive distribution into the tissues. Celecoxib is not preferentially bound to red blood cells.

Metabolism

Celecoxib metabolism is primarily mediated by CYP2C9. Three metabolites, a primary alcohol, the corresponding carboxylic acid and its glucuronide conjugate, have been identified in human plasma. These metabolites are inactive as COX-1 or COX-2 inhibitors.

Excretion

Celecoxib is eliminated predominantly by hepatic metabolism with little (<4%) unchanged drug recovered in the urine and feces. Following a single oral dose of radiolabeled drug, approximately 57% of the dose was excreted in the feces and 27% was excreted into the urine. The primary metabolite in both urine and feces was the carboxylic acid metabolite (73% of dose) with low amounts of the glucuronide also appearing in the urine. It appears that the low solubility of the drug prolongs the absorption process making terminal half-life (t_{1/2}) determinations more variable. The effective half-life is approximately 11 hours under fasted conditions. The apparent plasma clearance (CL/F) is about 500 mL/min.

Geriatric

At steady state, elderly subjects (over 65 years old) had a 40% higher C_{max} and a 50% higher AUC compared to the young subjects. In elderly females, celecoxib C_{max} and AUC are higher than those for elderly males, but these increases are predominantly due to lower body weight in elderly females. Dose adjustment in the elderly is not generally necessary. However, for patients of less than 50 kg in body weight, initiate therapy at the lowest recommended dose [see *DOSEAGE AND ADMINISTRATION* (2.6) and *USE IN SPECIFIC POPULATIONS* (8.3)].

Pediatric

The steady state pharmacokinetics of celecoxib administered as an investigational oral suspension was evaluated in 152 JRA patients 2 years to 17 years of age weighing 2 to 100 kg with psoriatic/arthritis or polyarthralgia course JRA and in patients with systemic onset JRA. Population pharmacokinetic analysis indicated that the oral clearance (adjusted for body weight) of celecoxib increases less than proportionally to increasing weight, with 10 kg and 25 kg patients predicted to have 40% and 24% lower clearance, respectively, compared to a 70 kg patient. The half-life of celecoxib was 11.2 hours in patients 2 to 17 years of age.

Twice-daily administration of 50 mg capsules to JRA patients weighing 212 to ≤ 25 kg and 100 mg capsules to JRA patients weighing >25 kg should achieve plasma concentrations similar to those observed in a clinical trial that demonstrated the non-inferiority of celecoxib to naproxen 7.5 mg/kg twice daily [see *DOSEAGE AND ADMINISTRATION* (2.3)]. Celecoxib has not been studied in JRA patients under the age of 2 years, in patients with body weight less than 10 kg (22 lbs), or beyond 24 weeks.

Race

Meta-analysis of pharmacokinetic studies has suggested an approximately 40% higher AUC of celecoxib in Blacks compared to Caucasians. The cause and clinical significance of this finding is unknown.

Hepatic Insufficiency

A pharmacokinetic study in subjects with mild (Child-Pugh Class A) and moderate (Child-Pugh Class B) hepatic impairment has shown that steady-state celecoxib AUC is increased about 40% and 180%, respectively, above that seen in healthy control subjects. Therefore, the daily recommended dose of celecoxib capsules should be reduced by 50% in patients with moderate (Child-Pugh Class B) hepatic impairment. Patients with severe hepatic impairment (Child-Pugh Class C) have not been studied. The use of celecoxib in patients with severe hepatic impairment is not recommended [see *DOSEAGE AND ADMINISTRATION* (2.6) and *USE IN SPECIFIC POPULATIONS* (8.6)].

Renal Insufficiency

In a cross-study comparison, celecoxib AUC was approximately 40% lower in patients with chronic renal insufficiency (GFR 35 to 60 mL/min) than that seen in subjects with normal renal function. No significant relationship was found between GFR and celecoxib clearance. Patients with severe renal insufficiency have not been studied. Similar to other NSAIDs, celecoxib is not recommended in patients with severe renal insufficiency [see *WARNINGS AND PRECAUTIONS* (8.6)].

Drug Interactions

In vitro studies indicate that celecoxib is not an inhibitor of cytochrome P450 2C9, 2C19 or 3A4. *In vivo* studies have shown the following:

Lithium: In a study conducted in healthy subjects, mean steady-state lithium plasma levels increased approximately 17% in subjects receiving lithium 450 mg twice daily with celecoxib 200 mg twice daily as compared to subjects receiving lithium alone [see *DRUG INTERACTIONS* (7.2)].

Fluconazole: Concomitant administration of fluconazole at 200 mg once daily resulted in a two-fold increase in celecoxib plasma concentration. This increase is due to the inhibition of celecoxib metabolism via P450 2C9 by fluconazole [see *DRUG INTERACTIONS* (7.5)].

Other Drugs: The effects of celecoxib on the pharmacokinetics and/or pharmacodynamics of glyburide, ketconazole, methotrexate [see *DRUG INTERACTIONS* (7.7)], phenytoin, and tolbutamide have been studied *in vivo* and clinically important interactions have not been found.

12.5 Pharmacogenomics

CYP2C9 activity is reduced in individuals with genetic polymorphisms that lead to reduced enzyme activity, such as those homozygous for the CYP2C9*2 and CYP2C9*3 polymorphisms. Limited data from 4 published reports that included a total of 8 subjects with the homozygous CYP2C9*3/*3 genotype showed celecoxib systemic levels that were 3- to 7-fold higher in these subjects compared to subjects with CYP2C9*1/*1 or *1/*3 genotypes. The pharmacokinetics of celecoxib have not been evaluated in subjects with other CYP2C9 polymorphisms, such as *2, *5, *6, *9 and *11. It is estimated that the frequency of the homozygous *3/*3 genotype is 0.3% to 1.0% in various ethnic groups. [see *DOSEAGE AND ADMINISTRATION* (2.6), *USE IN SPECIFIC POPULATIONS* (8.3)].

13. NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Celecoxib was not carcinogenic in rats given oral doses up to 200 mg/kg for males and 10 mg/kg for females (approximately 2 to 4 fold the human exposure as measured by the AUC₀₋₁₂ at 200 mg twice daily) or in mice given oral doses up to 25 mg/kg for males and 50 mg/kg for females (approximately equal to human exposure as measured by the AUC₀₋₁₂ at 200 mg twice daily) for two years.

Celecoxib was not mutagenic in an Ames test and a mutation assay in Chinese hamster ovary (CHO) cells, nor clastogenic in a chromosome aberration assay in CHO cells and an *in vivo* micronucleus test in rat bone marrow.

Celecoxib did not impair male fertility in rats at oral doses up to 600 mg/kg/day (approximately 11-fold human exposure at 200 mg twice daily based on the AUC₀₋₁₂).

13.2 Animal Toxicology

An increase in the incidence of background findings of spermatocleus with or without secondary changes such as epididymal hypoplasia, as well as an increase in the incidence of testicular atrophy and/or hypospadias seen in the juvenile rat. These reproductive findings while apparently treatment-related did not increase in incidence or severity with dose and may indicate an exacerbation of a spontaneous condition. Similar reproductive findings were not observed in studies of juvenile or adult dogs or in adult rats treated with celecoxib. The clinical significance of this observation is unknown.

14. CLINICAL STUDIES

14.1 Osteoarthritis

Celecoxib has demonstrated significant reduction in joint pain compared to placebo. Celecoxib was evaluated for treatment of the signs and the symptoms of OA of the knee and hip in placebo- and active-controlled clinical trials of up to 24 weeks in duration. In patients with OA, treatment with celecoxib 100 mg twice daily or celecoxib 200 mg once daily resulted in improvement in WOMAC (Western Ontario and McMaster Universities) osteoarthritis index, a composite of pain, stiffness, and functional measures in OA. In three 12-week studies of pain accompanying OA flare, celecoxib doses of 100 mg twice daily and 200 mg twice daily provided significant reduction of pain within 24 to 48 hours of initiation of dosing. At doses of 100 mg twice daily or 200 mg twice daily the effectiveness of celecoxib was shown to be similar to that of naproxen 500 mg twice daily. Doses of 200 mg twice daily provided no additional benefit above that seen with 100 mg twice daily. A total daily dose of 200 mg has been shown to be equally effective whether administered as 100 mg twice daily or 200 mg once daily.

14.2 Rheumatoid Arthritis

Celecoxib has demonstrated significant reduction in joint tenderness/pain and joint swelling compared to placebo. Celecoxib was evaluated for treatment of the signs and symptoms of JRA in placebo- and active-controlled clinical trials of up to 24 weeks