

Trivagizole 3TM

Clotrimazole Vaginal Cream (2%)
Vaginal Antifungal

Vaginal Cream

3 Day Treatment

1 Reusable Applicator

Trivagizole 3TM

Clotrimazole Vaginal Cream (2%)
Vaginal Antifungal

Vaginal Cream

Cures Most Vaginal Yeast Infections

Trivagizole 3TM

Clotrimazole Vaginal Cream (2%)
Vaginal Antifungal

Vaginal Cream

One 25 g (0.88 oz) Tube Of Clotrimazole Vaginal Cream (2%)
and 1 Reusable Applicator

Cures Most
Vaginal Yeast
Infections

3 Day Treatment



Educational Brochure Enclosed



Drug Facts (continued)

Other information

- Store at 20-25°C (68-77°F)
- See end flap of carton and tube for lot number and expiration date
- The tube opening should be sealed. If seal has been broken, do not use product.

Inactive ingredients benzyl alcohol, purified water, sorbitan monostearate, canol, polyoxamate-60, purified water, sorbitan monostearate

Questions or comments? Call toll-free, 1-888-827-8222, for our 24 hour automated response system.

TARO

TARO

TARO

Distributed by:
Taro Pharmaceuticals
U.S.A., Inc.
Hawthorne, NY 10532
Taro is a registered trademark of
Taro Pharmaceuticals U.S.A., Inc.
www.taropharma.com



Drug Facts

Active ingredient (in each applicator)
Clotrimazole 100 mg (2%)
Vaginal Antifungal

Use
Treats vaginal yeast infections

Warnings
For vaginal use only

Do not use
■ If you have never had a vaginal yeast infection diagnosed by a doctor
■ If you have a fever (higher than 100°F), pain in the lower abdomen, back, or other shoulder, or a foul-smelling vaginal discharge. You should see a doctor for treatment of these symptoms.

Ask a doctor before use if you have
■ vaginal itching and discomfort for the first time
■ symptoms that return within 2 months. You could be pregnant, or have a serious underlying medical cause for your symptoms, including diabetes or a weakened immune system (which may be due to HIV - the virus that causes AIDS).

When using this product
■ do not use tampons, douches, or spermicides
■ do not use condoms or diaphragms, as this product may damage them and they may not prevent sexually transmitted disease or pregnancy

Stop use and ask a doctor if
■ symptoms do not get better in 3 days
■ you get a rash, fever, abdominal pain, or a foul-smelling vaginal discharge

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions
■ Before using, read and keep the enclosed educational brochure for complete instructions and information
■ adults and children 12 years and over:
■ insert one applicatorful of cream into the vagina at bedtime for 3 days in a row
■ children under 12 years: ask a doctor

3 Day Treatment
1 Reusable Applicator
Cures Most Vaginal Yeast Infections

APPROVAL		
DATE	3/22/00	ANT COPY CTRL.0

CAP END

127 mm
TUBE
LENGTH

OPEN END

TARO

Trivagizole³™

Clotrimazole Vaginal Cream (2%)
Vaginal Antifungal

3 Day Treatment

Weight 25g (0.88oz)

For vaginal use only. Do not use in eyes or take by mouth.
If this is the first time you have had vaginal itching and discomfort, consult your doctor. If you have had a doctor diagnose a vaginal yeast infection before and have the same symptoms now, use this cream as directed for 3 days in a row.
Warnings: Read the Warnings and Directions on the carton and in the educational brochure before using this product.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.
Directions: To Open: Unscrew the cap, use pointed end on cap to break seal. Fill the applicator and insert one applicatorful of cream into vagina at bedtime for 3 days in a row.
Active Ingredient: Clotrimazole 2% (100 mg per applicator)
Store at 20-25 °C (68-77 °F).
See end of tube for lot number and expiration date.
Safety Sealed: The tube opening should be sealed. If seal has been broken, do not use product.
Mfd. by: Taro Pharmaceuticals Inc.
Bramalea, Ontario, Canada L6T 1C3 Made in Canada PK-0000-0

TARO

PMS 281

PK-0000-0

APPROVAL					
DATE 3/17/00		LPS		ART COPY CTRL.#	
PASS		UPC		PHARMACODE	
DEPARTMENT	SIGNATURE	NO	YES	YES W CORR	DATE
PROOFING					
ART					
MARKETING					
ENGINEERING					
O.C.					
REG.					
CLIENT					
THIS COLOR PROOF INDICATES COLOR BREAK ONLY AND MAY NOT ACCURATELY REFLECT ACTUAL PRODUCTION COLOR. IF YOU HAVE ANY QUESTIONS OR COMMENTS PLEASE CALL TARO PHARMACEUTICALS-ART DEPARTMENT 800-844-1448.					

ID BAR CODE

EDUCATIONAL BROCHURE

Read This Before Using

Trivagizole 3™

Clotrimazole

Vaginal

Cream (2%)

Vaginal Antifungal

Contains 2% (100 mg per applicator)

3-day treatment of vaginal yeast infections.

Use:

• For the treatment of vaginal yeast infections.

If you have symptoms such as vaginal itching, burning, or discharge, and if at some time in the past your doctor has told you that these symptoms were due to a vaginal yeast infection, then Trivagizole 3 vaginal Cream should work for you. If, however, you have never had these symptoms or a vaginal yeast infection before, you should see your doctor before using this product.

What is a Vaginal Yeast Infection?

A yeast infection is a common type of vaginal infection. Your doctor may call it candidiasis. This condition is caused by an

organism called *Candida*, which is a type of yeast.

Healthy women usually have this yeast on the skin, in the mouth, in the digestive tract, and in the vagina. At times, the yeast can grow very quickly. In fact, the infection is sometimes called yeast (*Candida*) "overgrowth." Some women may have a yeast infection on the skin outside the vagina (vulva), associated with the vaginal infection.

A yeast infection can occur at any age. It is most common during the childbearing years. Women who are pregnant or diabetic, taking antibiotics, birth control pills or steroids, or who have a weakened immune system are more likely to get repeated yeast infections.

Various medical conditions can damage the body's normal defenses against

infection. One of the most serious of these conditions is infection with the human immunodeficiency virus (HIV), the virus that causes AIDS. The HIV virus causes the body to be more likely to get infections, including vaginal yeast infections.

Therefore, women with HIV infection may have frequent vaginal yeast infections that do not clear up easily with proper treatment. If you may have been exposed to HIV and are experiencing repeated vaginal yeast infections, you should see your doctor right away for additional information on HIV infection, please contact your doctor or the CDC National AIDS Hotline. The CDC phone numbers are: 1-800-342-AIDS (English), 1-800-344-7432 (Spanish), or 1-800-243-7889 (hearing impaired, TDD).

Symptoms of a Vaginal Yeast Infection

There are many symptoms of a vaginal yeast infection. They can include:

- Vaginal itching (longing from mild to severe).
- A clumpy, vaginal discharge that may look like cottage cheese.
- Vaginal soreness, irritation, or burning.
- Itchiness on the skin outside the vagina (vulvar irritation).

Note: Vaginal discharge that is different from above, for example, a yellow/green discharge or a foul-smelling discharge may mean that you have a different infection. If this is the case, you should talk to your doctor before using Trivagizole 3 vaginal Cream.

Warnings

• For vaginal use only. Do not use in eyes or take by mouth.

• This product is effective only in treating vaginal yeast infections.

• Do not use Trivagizole 3 Vaginal Cream if you have any of the following symptoms:

- Fever (higher than 100°F).
- Pain in the lower abdomen, back, or either shoulder.
- A foul-smelling vaginal discharge.

• While using Trivagizole 3 Vaginal Cream, if you get a fever, abdominal pain, or a foul-smelling vaginal discharge, stop

using the product & contact your doctor at once. You may have more serious disease.

• If your symptoms do not improve in 3 days, you still have symptoms after 7 days, you should call your doctor.

• If your symptoms (even within 2 months) should talk to your doctor. You could be pregnant or there could be a serious underlying medical cause for your infection, including diabetes or a weakened immune system (which may be due to HIV, the virus that causes AIDS).

• Contact your doctor if you get a fever, abdominal pain, or a foul-smelling vaginal discharge while using this product.

4-10 CLOTRIMAZOLE 2% CR, 3-DAY (TARO) - INSERT - FRONT 4 X 12 FOLDED TO 1 X DIE IS AT 100%

COLORS...
PMS-291 Date 3/21/00

page 4

Signs and Symptoms	Warnings	Using the product and contact your doctor right away. You may have a more serious disease.	Do not use tampons, douches, or spermicides while using this medicine.
• (tongue, throat, or mouth) infection	• This product is effective only in treating vaginal yeast infections.	• If your symptoms do not improve in 3 days, or you still have symptoms after 7 days, you should call your doctor.	• Do not rely on condoms or diaphragms to prevent sexually transmitted diseases or pregnancy. This product may damage condoms and diaphragms and cause them to fail.
• Do not use Trivagizole 3 Vaginal Cream if you have any of the following symptoms:	• Do not use Trivagizole 3 Vaginal Cream if you have any of the following symptoms:	• If your symptoms return within 2 months, you should talk to your doctor. You could be pregnant or there could be a serious underlying medical cause for your infections, including diabetes or a weakened immune system (which may be due to HIV - the virus that causes AIDS).	• If pregnant or breastfeeding, ask a health professional before use.
• Itching that does not go away after you wash the area.	• Pain in the lower abdomen, back, or either shoulder.	• Contact your doctor if you get hives or a skin rash while using this product.	• Do not use in girls less than 12 years of age.
• A foul-smelling vaginal discharge.	• While using Trivagizole 3 Vaginal Cream, if you get a fever, abdominal pain, or a foul-smelling vaginal discharge, stop using the product.		• Keep this and all drugs out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Boxes print
8% BD
PMS 281

INSERT - FRONT 4 X 12 FOLDED TO 1 X 1/2

COLORS...
PMS 291
Date 3/21/00

Side Effects

A mild increase in vaginal burning, itching, or irritation may occur when Trivagazole 3 Vaginal Cream is inserted. Stop using Trivagazole 3 Vaginal Cream and tell your doctor if you have abdominal pain, hives, or skin rash, or if you have severe vaginal burning, itching, or irritation.

For Best Results

1. Be sure to use this product for the full 3 days, even if your symptoms go away sooner.
2. Use one applicatorful of Trivagazole 3 Vaginal Cream at bedtime for three days in a row, even during your menstrual period.
3. Wear cotton underwear.
4. If your partner has itching, redness, or discomfort of his penis, he should consult his doctor. He should tell the doctor that you are treating

Yoursell for a vaginal yeast infection

5. Dry the outside vaginal area thoroughly, after a shower, bath, or swim.
6. Change out of a wet bathing suit or damp clothes as soon as possible. A dry area is less likely to lead to the growth of yeast.
7. Wipe from front to rear after urination.
8. Do not douche unless your doctor tells you to do so. Douching may wash the product out of the vagina.
9. Try not to scratch the skin outside the vagina. Scratching can cause more irritation and can spread the infection.
10. Discuss with your doctor any drugs you are now taking. You are more likely to get a vaginal yeast infection with certain drugs.
11. Do not have vaginal intercourse while using this product.

If you have a question

Questions of a medical nature should be taken up with your doctor. If you have any other questions, or need more information on this product, call our 24 hour automated response system, TOLL-FREE at 1-888-827-6222, Monday through Friday.

Active ingredient:

Clotrimazole 2% (100 mg per applicator)

Inert ingredients:

Benzyl alcohol, cetostearyl alcohol, cetyl esters wax, 2-octyldodecanol, polysorbate 80, purified water, sorbitol monosorbate

Storage

Store at 20-25°C (68-77°F). See end flap of carton and end of tube for lot number and expiration date.

How to use Trivagazole 3 Vaginal Cream with 1 Reusable Applicator


Directions for Use

Use the cream of bedtime for 3 days in a row. Read the full directions below before using.


1. To open the tube, unstick the cap.




Press the sharp point of the cap into the sealed end of the tube. Push down firmly until the seal is broken (see picture).




2. Attach the applicator to the tube of cream by placing end of the applicator firmly onto tube of cream (see picture).




3. Gently squeeze the cream into the applicator. Continue squeezing until line on plunger is at FULL. Leeches and oil barrel. Separate the applicator from the tube. After each use, replace cap and roll up the tube from the bottom.



4. Fold the label as shown in the picture.



5. The Trivagazole 3 Vaginal Cream applicator is inserted into the vagina in much the same way as a tampon. Gently insert the applicator into the vagina as far back as it will go comfortably. This can be done while lying on your back with your knees bent (see picture), or while sitting with your feet open and knees bent.



6. With one hand, hold your back with your bent (see picture), or while sitting with your feet open and knees bent.



7. Gently use the other hand to push the plunger down for as far as it will go. Remove the applicator from the vagina.

FPI


4-10 CLOTRIMAZOLE 2% CR, 3-DAY (TARP) - INSERT - BACK VIEW 4 X 12 FOLDED TO 1 X
DIE IS AT 100%

COLORS...
PMS 287 Date 3/17/00


page 6

ID BAR CODE


2. Attach the applicator to the tube of cream by placing end of the cap, actor firmly onto the of cream. (See picture.)




4. Hold the barrel as shown in the picture.



5. The Tivrogzole 3 Vaginal Cream applicator is inserted into the vagina in much the same way as a tampon. Gently insert the applicator into the vagina as far back as it will go comfortably. This can be done while lying on



your back with your knees bent. (As shown in the picture), or while standing with your feet apart and your knees bent.



and warm water. Rinse well and dry. Put the two pieces back together and store in a clean, dry place.

8. Lie down as soon as possible to keep the cream from leaking out.

9. Repeat steps 2 through 8 before bedtime for the next 2 days.

To protect your clothing from leakage or discharge, you may want to use deodorant, free pads or panty shields while you are using Tivrogzole 3 Vaginal Cream. Do not use tampons, douches, or spermicides until after you have completed the treatment and your symptoms are gone.

Mfg. by:
Tara Pharmaceuticals Inc.
Bromfield, Ontario
Canada L6T 1C3
www.tivrogzole.com

FPI

barrel, use the other hand to push the plunger into the barrel as far as it will go. Remove the applicator from the vagina.

7. Clean the applicator after each use. Pull the two pieces apart. Wash them with soap

3. Gently squeeze the cream into the applicator. Continue squeezing until the cream is fully loaded. If ULL reaches end of barrel. Separate the applicator from the tube. After each use, replace cap and roll up the tube from the bottom.

NO - INSERT - BACK VIEW

4 X 12 FOLDED TO 1 X 1/2

COLORS...

PMS 661

Date 3/17/00