



PAA176915.FDA02



FRONT




BACK

AREA FOR LOT AND EXP
TO BE PRINTED IN DARK COLOR

PAA117156.FDA02

MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• BARLINES, HAIRLINES	N/A

FRONT



BACK



AREA FOR LOT AND EXP
TO BE PRINTED IN DARK COLOR
26 mm x 5.5 mm

PAA117157.FDA02

MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• BARLINES, HAIRLINES	N/A



PAA117160.FDA02

NON MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	N/A
• DRUG FACTS CONTINUED	N/A
• HEADINGS	N/A
• SUBHEADINGS/BODY TEXT	N/A
• LEADING	N/A
• # OF CHARACTERS PER INCH	N/A
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• BARLINES, HAIRLINES	1pt
• SPACE BETWEEN HAIRLINES AND BOX END	N/A

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Do not use with other medicines containing ACETAMINOPHEN; can cause liver damage.

Drug Facts

Active ingredients (in each caplet)

Purposes

Acetaminophen 250 mgPain reliever

Ibuprofen 125 mg (NSAID*)Pain reliever

*nonsteroidal anti-inflammatory drug

Uses

■ temporarily relieves minor aches and pains due to:

■ headache

■ toothache

■ backache

■ menstrual cramps

■ muscular aches

■ minor pain of arthritis

PA118118.FDA02

LIFT HERE

PANEL 1

Drug Facts (continued)

Warnings

Acetaminophen liver damage warning:

This product contains acetaminophen. Severe liver damage may occur if you take:

■ with other drugs containing acetaminophen

■ more than 6 caplets in 24 hours, which is the maximum daily amount for this product

■ 3 or more alcoholic drinks every day while using this product

Acetaminophen allergy alert: may cause severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash

If skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

NSAID allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

■ hives

■ facial swelling

■ asthma (wheezing)

■ shock

■ skin reddening

■ rash

■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

NSAID stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

■ are age 60 or older

■ have had stomach ulcers or bleeding problems

PANEL 2

Drug Facts (continued)

■ take a blood thinning (anticoagulant) or steroid drug

■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

■ have 3 or more alcoholic drinks every day while using this product

■ take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription).

Drug Facts (continued)

If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

■ if you have ever had an allergic reaction to acetaminophen or any other pain reliever

■ right before or after heart surgery

Ask a doctor before use if

■ you have liver disease

■ stomach bleeding warning applies to you

■ you have problems or serious side effects from taking pain relievers

■ you have a history of stomach problems, such as heartburn

■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

■ you are taking a diuretic

PANEL 3

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

■ under a doctor's care for any serious condition

■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin

■ taking any other drug

When using this product

■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding:

■ feel faint

■ vomit blood

■ have bloody or black stools

■ have stomach pain that does not get better

Drug Facts (continued)

■ you have symptoms of heart problems or stroke:

■ chest pain

■ trouble breathing

■ weakness in one part or side of body

■ slurred speech

■ leg swelling

■ pain gets worse or lasts more than 10 days

■ redness or swelling is present in the painful area

■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

PANEL 4

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ do not take more than directed

adults and children 12 years and over

■ take 2 caplets every 8 hours while symptoms persist

children under 12 years

■ ask a doctor

■ do not take more than 6 caplets in 24 hours, unless directed by a doctor

Other information

■ read all warnings and directions before use. Keep carton.

■ store at 20-25°C (68-77°F)

■ avoid excessive heat above 40°C (104°F)

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, ferric oxides, glyceryl dibehenate, hypromellose, pharmaceutical ink, polydextrose, polyethylene glycol, pregelatinized starch, titanium dioxide

Questions or comments?

call weekdays 9 AM to 5 PM at 1-800-88-ADVIL

BASE PANEL

PAA118118.FDA02

NON MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

ADVIL® SAFETY SEALED ADVIL® SAFETY SEALED ADVIL® SAFETY SEALED

TEAR HERE TO OPEN

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

■ under a doctor's care for any serious condition

■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin

■ taking any other drug

When using this product

■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding:

■ feel faint

■ vomit blood

■ have bloody or black stools

■ have stomach pain that does not get better

■ you have symptoms of heart problems or stroke:

■ chest pain

■ trouble breathing

■ weakness in one part or side of body

■ slurred speech

■ leg swelling

■ pain gets worse or lasts more than 10 days

■ redness or swelling is present in the painful area

■ any new symptoms appear

If pregnant or breast-feeding

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children

. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ do not take more than directed

adults and children 12 years and over

■ take 2 caplets every 8 hours while symptoms persist

children under 12 years

■ ask a doctor

■ do not take more than 6 caplets in 24 hours, unless directed by a doctor

Other information

■ read all warnings and directions before use.

■ store at 20-25°C (68-77°F)

■ avoid excessive heat above 40°C (104°F)

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, ferric oxides, glyceryl dibehenate, hypromellose, pharmaceutical ink, polydextrose, polyethylene glycol, pregelatinized starch, titanium dioxide

Questions or comments?

call weekdays 9 AM to 5 PM at 1-800-88-ADVIL

For most recent product information, visit [www.Advil.com](#)

For US Patent or Application status see [www.productpats.com](#)

Pfizer, Madison, NJ 07940 USA © 2020 Pfizer Inc.

PAA117158.FDA03

81% For Position Only

3 0573 0147 09 5

CONTAINS 2 MEDICINES

Advil®

DUAL ACTION

WITH **Acetaminophen**

Acetaminophen 250 mg and Ibuprofen (NSAID) 125 mg Tablets

Pain Reliever

8 Caplets*

*Capsule-Shaped Tablets

DO NOT USE IF SEAL AROUND CAP IS BROKEN OR MISSING. ADVIL SAFETY SEALED.

Do not use with other medicines containing ACETAMINOPHEN; can cause liver damage.

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg

Pain reliever

Ibuprofen 125 mg (NSAID*)

Pain reliever

*nonsteroidal anti-inflammatory drug

Purposes

LIFT HERE for More Drug Facts

UNVARNISHED AREA FOR LOT & EXP. DATE

Drug Facts (continued)

Uses

■ temporarily relieves minor aches and pains due to:

■ headache

■ toothache

■ backache

■ menstrual cramps

■ muscular aches

■ minor pain of arthritis

Warnings

Acetaminophen liver damage warning:

This product contains acetaminophen. Severe liver damage may occur if you take:

■ with other drugs containing acetaminophen

■ more than 6 caplets in 24 hours, which is the maximum daily amount for this product

■ 3 or more alcoholic drinks every day while using this product

Acetaminophen allergy alert:

may cause severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash

If skin reaction occurs, stop use and seek medical help right away.

NSAID allergy alert:

ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

■ hives

■ facial swelling

■ asthma (wheezing)

■ shock

■ skin reddening

■ rash

■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

NSAID stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

■ are age 60 or older

■ have had stomach ulcers or bleeding problems

■ take a blood thinning (anticoagulant) or steroid drug

■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

■ have 3 or more alcoholic drinks every day while using this product

■ take more or for a longer time than directed

Heart attack and stroke warning:

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

■ if you have ever had an allergic reaction to acetaminophen or any other pain reliever

■ right before or after heart surgery

Ask a doctor before use if

■ you have liver disease

■ stomach bleeding warning applies to you

■ you have problems or serious side effects from taking pain relievers.

■ you have a history of stomach problems, such as heartburn

■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke



■ you are taking a diuretic

PAA117158.FDA03

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

1.25" X 0.5" LOT & EXP BOX.
NO TEXT AREA AND
PRINTED IN A DARK COLOR.





PAA117201.FDA02

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	6 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	.5 pt

TOP (BACK)

Warnings**Acetaminophen liver damage warning:**

This product contains acetaminophen. Severe liver damage may occur if you take: with other drugs containing acetaminophen; more than 6 caplets in 24 hours, which is the maximum daily amount for this product; 3 or more alcoholic drinks every day while using this product.

Acetaminophen allergy alert: may cause severe skin reactions.

Symptoms may include: skin reddening, blisters, and rash. If skin reaction occurs, stop use and seek medical help right away.

NSAID allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Do not use this product if you have ever had an allergic reaction to any pain reliever; with any drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist; or right before or after surgery.

NSAID stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinning (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed.

BASE

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Ask a doctor before use if you are pregnant or breast feeding: you have liver disease; stomach bleeding warning applies to you; you have problems or serious side effects from taking pain relievers; you have a history of stomach problems, such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; you are taking a diuretic.

Ask your doctor or pharmacist before use if you are under a doctor's care for any serious condition; taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin; or taking any other drug.

When using this product take with food or milk if stomach upset occurs.

Stop use and ask a doctor if pain gets worse or lasts more than 10 days; redness or swelling is present in the painful area; or any new symptoms appear.

Directions: do not take more than directed; adults and children 12 years and over: **take 2 caplets every 8 hours** while symptoms persist; do not take more than 6 caplets in 24 hours, unless directed by a doctor; children under 12 years: ask a doctor.

(Questions or comments? 1-800-88-ADVIL)

DO NOT USE unless vial was sealed in unbroken plastic blister on cardboard card.

For most recent product information, visit www.Advii.com
Pfizer, Madison, NJ 07940 USA © 2020 Pfizer Inc.

PAA116916.FDA02

TOP (FACE)

CONTAINS 2 MEDICINES

Advil®
DUAL ACTION
WITH **Acetaminophen**

Acetaminophen 250 mg and
Ibuprofen (NSAID) 125 mg Tablets
Pain Reliever

Advil II 8 Caplets

READ AND KEEP CARD FOR COMPLETE WARNINGS AND INFORMATION

Uses
temporarily relieves minor aches and pains due to:



- headache
- backache
- muscular aches
- toothache
- menstrual cramps
- minor pain of arthritis

OPEN HERE TO VIEW MORE PRODUCT INFORMATION

LOT EXP AREA
NO VARNISH

PAA116916.FDA02


DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	6 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE	.5 pt




PAA117202.FDA02

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

FRONT



BACK



LOT & EXP AREA

PAA117009.FDA02

DRUG FACTS TEXT DEFINED	TYPE SIZE
• TEXT	6 pt
• LEADING	6.5 pt
• BARLINES, HAIRLINES	.75 pt

PANEL 2

Drug Facts (continued)

NSAID allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

NSAID stomach bleeding warning:
This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you have ever had an allergic reaction to acetaminophen or any other pain reliever
- right before or after heart surgery

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers
- you have a history of stomach problems, such as heartburn

Drug Facts (continued)

- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

PANEL 3

PANEL 4

Drug Facts (continued)

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days

Drug Facts (continued)

- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts

Active ingredients (in each caplet)	Purposes
Acetaminophen 250 mg Pain reliever	
Ibuprofen 125 mg (NSAID*) Pain reliever *nonsteroidal anti-inflammatory drug	

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - muscular aches
 - minor pain of arthritis

Drug Facts (continued)

Warnings
Acetaminophen liver damage warning:
This product contains acetaminophen. Severe liver damage may occur if you take:

- with other drugs containing acetaminophen
- more than 6 caplets in 24 hours, which is the maximum daily amount for this product
- 3 or more alcoholic drinks every day while using this product

Acetaminophen allergy alert: may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If skin reaction occurs, stop use and seek medical help right away.

BASE PANEL

Drug Facts (continued)

Directions

- do not take more than directed

adults and children 12 years and over	■ take 2 caplets every 8 hours while symptoms persist
children under 12 years	■ ask a doctor

- do not take more than 6 caplets in 24 hours, unless directed by a doctor

Other information

- read all warnings and directions before use.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)


Inactive ingredients carnauba wax, colloidal silicon dioxide, croscarmellose sodium, ferric oxides, glyceryl dibehenate, hypromellose, pharmaceutical ink, polydextrose, polyethylene glycol, pregelatinized starch, titanium dioxide

Questions or comments? call weekdays 9 AM to 5 PM at 1-800-88-ADVIL

PAA118119.FDA02

PAA118119.FDA02

NON MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



PAA116859.FDA02

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

TOP (FACE)
PANEL 1

CONTAINS 2 MEDICINES

Advil®
DUAL ACTION
with **Acetaminophen**

Acetaminophen 250 mg and
Ibuprofen (NSAID) 125 mg Tablets
Pain Reliever

18
Caplets

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Uses temporarily relieves minor aches and pains due to: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ muscular aches ■ minor pain of arthritis

Warnings

Acetaminophen liver damage warning: This product contains acetaminophen. Severe liver damage may occur if you take: with other drugs containing acetaminophen;

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing, more than 6 caplets in 24 hours, which is the maximum daily amount for this product; 3 or more alcoholic drinks every day while using this product.

Acetaminophen allergy alert: may cause severe skin reactions. Symptoms may include: skin reddening, blisters, and rash. If skin reaction occurs, stop use and seek medical help right away.

PAA116860.FDA02 **LIFT HERE for More Drug Facts**

FOR LOT & EXPIRATION AREA,
UNVARNISHED AND PRINTED
IN BLACK OR DARK BLUE.

BASE
PANEL 3

heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; you are taking a diuretic.

Ask your doctor or pharmacist before use if you are under a doctor's care for any serious condition; taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin; or taking any other drug.

When using this product take with food or milk if stomach upset occurs.

Stop use and ask a doctor if pain gets worse or lasts more than 10 days; redness or swelling is present in the painful area; or any new symptoms appear.

Directions

■ **do not take more than directed**

adults and children 12 years and over	■ take 2 caplets every 8 hours while symptoms persist
children under 12 years	■ ask a doctor

■ **do not take more than 6 caplets in 24 hours, unless directed by a doctor**

Questions or comments? 1-800-88-ADVIL
Pfizer, Madison, NJ 07940 USA © 2020 Pfizer Inc.
For most recent product information, visit www.Advil.com

TOP (BACK)
PANEL 2

NSAID allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Do not use this product if you have ever had an allergic reaction to any pain reliever; with any drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist; or right before or after surgery.

NSAID stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers

or bleeding problems; take a blood thinning (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]; have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Ask a doctor before use if you are pregnant or breast feeding; you have liver disease; stomach bleeding warning applies to you; you have problems or serious side effects from taking pain relievers; you have a history of stomach problems, such as heartburn; you have high blood pressure,

PAA116860.FDA02

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	6 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE	.5 pt



PAA116918.FDA02


DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

TOP (FACE) PANEL 1




FOR LOT & EXPIRATION AREA,
UNVARNISHED AND PRINTED
IN BLACK OR DARK BLUE.

BASE PANEL 3




TOP (BACK) PANEL 2



PAA118181.FDA02

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	6 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE	.5 pt



DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

TOP (FACE)
PANEL 1

CONTAINS 2 MEDICINES

Advil®

DUAL ACTION

with **Acetaminophen**

Acetaminophen 250 mg and
Ibuprofen (NSAID) 125 mg Tablets
Pain Reliever

Advil

36
Caplets

READ AND KEEP CARTON FOR COMPLETE
WARNINGS AND INFORMATION

Uses temporarily relieves minor aches and pains due to: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ muscular aches ■ minor pain of arthritis

Warnings

Acetaminophen liver damage warning: This product contains acetaminophen. Severe liver damage may occur if you take: with other drugs containing acetaminophen;

Do Not Use if seal under bottle cap imprinted with **"SEALED for YOUR PROTECTION"** is broken or missing. more than 6 caplets in 24 hours, which is the maximum daily amount for this product; 3 or more alcoholic drinks every day while using this product.

Acetaminophen allergy alert: may cause severe skin reactions. Symptoms may include: skin reddening, blisters, and rash. If skin reaction occurs, stop use and seek medical help right away.

PAA117206.FDA02 **LIFT HERE for More Drug Facts**

FOR LOT & EXPIRATION AREA,
UNVARNISHED AND PRINTED
IN BLACK OR DARK BLUE.

BASE
PANEL 3

heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; you are taking a diuretic.

Ask your doctor or pharmacist before use if you are under a doctor's care for any serious condition; taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin; or taking any other drug.

When using this product take with food or milk if stomach upset occurs.

Stop use and ask a doctor if pain gets worse or lasts more than 10 days; redness or swelling is present in the painful area; or any new symptoms appear.

Directions

■ **do not take more than directed**

adults and children 12 years and over	■ take 2 caplets every 8 hours while symptoms persist
children under 12 years	■ ask a doctor

■ **do not take more than 6 caplets in 24 hours, unless directed by a doctor**

Questions or comments? 1-800-88-ADVIL
Pfizer, Madison, NJ 07940 USA © 2020 Pfizer Inc.
For most recent product information, visit www.Advil.com

TOP (BACK)
PANEL 2

NSAID allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Do not use this product if you have ever had an allergic reaction to any pain reliever; with any drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist; or right before or after surgery.

NSAID stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers

or bleeding problems; take a blood thinning (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]; have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Ask a doctor before use if you are **pregnant or breast feeding**; you have liver disease; stomach bleeding warning applies to you; you have problems or serious side effects from taking pain relievers; you have a history of stomach problems, such as heartburn; you have high blood pressure,

PAA117206.FDA02

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	6 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE	.5 pt

Drug Facts (continued)

Other information

■ read all warnings and directions before use. Keep carton.

■ store at 20-25°C (68-77°F)

■ avoid excessive heat above 40°C (104°F)

Inactive ingredients carnauba wax, colloidal silicon dioxide, croscarmellose sodium, ferric oxides, glyceryl dibehenate, hypromellose, pharmaceutical ink, polydextrose, polyethylene glycol, pregelatinized starch, titanium dioxide

Questions or comments?
call weekdays 9 AM to 5 PM at 1-800-88-ADVIL

BASE PANEL

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

■ do not take more than directed

adults and children 12 years and over

■ take 2 caplets every 8 hours while symptoms persist

children under 12 years

■ ask a doctor

■ do not take more than 6 caplets in 24 hours, unless directed by a doctor

PANEL #4

Drug Facts (continued)

■ take a blood thinning (anticoagulant) or steroid drug

■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

■ have 3 or more alcoholic drinks every day while using this product

■ take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

11

LIFT HERE

PAA118116.FDA02

PANEL #1

Drug Facts (continued)

■ if you have ever had an allergic reaction to acetaminophen or any other pain reliever

■ right before or after heart surgery

Ask a doctor before use if

■ you have liver disease

■ stomach bleeding warning applies to you

■ you have problems or serious side effects from taking pain relievers

■ you have a history of stomach problems, such as heartburn

Drug Facts (continued)

■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are

■ under a doctor's care for any serious condition

■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin

■ taking any other drug

PANEL #2

Drug Facts (continued)

When using this product

■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding:

■ feel faint

■ vomit blood

■ have bloody or black stools

■ have stomach pain that does not get better

Drug Facts (continued)

■ you have symptoms of heart problems or stroke:

■ chest pain

■ trouble breathing

■ weakness in one part or side of body

■ slurred speech

■ leg swelling

■ pain gets worse or lasts more than 10 days

■ redness or swelling is present in the painful area

■ any new symptoms appear

PANEL #3

PAA118116.FDA02

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	N/A
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

CONTAINS 2 MEDICINES

Advil®

DUAL ACTION

WITH Acetaminophen

Acetaminophen 250 mg and Ibuprofen (NSAID) 125 mg Tablets Pain Reliever

Advil II

72 Caplets

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Uses temporarily relieves minor aches and pains due to:
■ backache ■ menstrual cramps ■ muscular aches ■ minor pain of arthritis

Warnings
Acetaminophen liver damage warning: This product contains acetaminophen. Severe liver damage may occur if you take with other drugs containing acetaminophen; more than 6 caplets in 24 hours, which is the maximum daily amount for this product; 3 or more alcoholic drinks every day while using this product.
Acetaminophen allergy alert: may cause severe skin reactions. Symptoms may include: skin reddening, blisters, and rash. If skin reaction occurs, stop use and seek medical help right away.
NSAID allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.
Do not use this product if you have ever had an allergic reaction to any pain reliever; with any drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist; or right before or after surgery.
NSAID stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinning (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed.
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Ask a doctor before use if you are pregnant or breast feeding; you have liver disease; stomach bleeding warning applies to you; you have problems or serious side effects from taking pain relievers; you have a history of stomach problems, such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; you are taking a diuretic.

Ask your doctor or pharmacist before use if you are under a doctor's care for any serious condition; taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin; or taking any other drug.
When using this product take with food or milk if stomach upset occurs.
Stop use and ask a doctor if pain gets worse or lasts more than 10 days; redness or swelling is present in the painful area; or any new symptoms appear.

Directions
■ do not take more than directed

adults and children 12 years and over	■ take 2 caplets every 6 hours while symptoms persist
children under 12 years	■ ask a doctor

■ do not take more than 6 caplets in 24 hours, unless directed by a doctor

Questions or comments? 1-800-98-ADVIL.
Pfizer, Madison, NJ 07940 USA © 2020 Pfizer Inc.
For most recent product information, visit www.Advii.com

PAA117205.FDA02

UNVARNISHED AND PRINTED IN BLACK OR DARK BLUE FOR LOT & EXPIRATION AREA.

PAA117205.FDA02

MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.95 pt
• BULLETS	3.5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	.5 pt



DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

CONTAINS 2 MEDICINES

Advil®

DUAL ACTION

WITH Acetaminophen

Acetaminophen 250 mg and Ibuprofen (NSAID) 125 mg Tablets Pain Reliever

Advil II

90 Caplets

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Uses temporarily relieves minor aches and pains due to:
■ backache ■ menstrual cramps ■ muscular aches ■ minor pain of arthritis

Warnings
Acetaminophen liver damage warning: This product contains acetaminophen. Severe liver damage may occur if you take with other drugs containing acetaminophen; more than 6 caplets in 24 hours, which is the maximum daily amount for this product; 3 or more alcoholic drinks every day while using this product.
Acetaminophen allergy alert: may cause severe skin reactions. Symptoms may include: skin reddening, blisters, and rash. If skin reaction occurs, stop use and seek medical help right away.
NSAID allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.
Do not use this product if you have ever had an allergic reaction to any pain reliever; with any drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist; or right before or after surgery.
NSAID stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinning (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed.
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Ask a doctor before use if you are pregnant or breast feeding; you have liver disease; stomach bleeding warning applies to you; you have problems or serious side effects from taking pain relievers; you have a history of stomach problems, such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; you are taking a diuretic.
Ask your doctor or pharmacist before use if you are under a doctor's care for any serious condition; taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin; or taking any other drug.
When using this product take with food or milk if stomach upset occurs.
Stop use and ask a doctor if pain gets worse or lasts more than 10 days; redness or swelling is present in the painful area; or any new symptoms appear.

Directions
■ do not take more than directed

adults and children 12 years and over	■ take 2 caplets every 6 hours while symptoms persist
children under 12 years	■ ask a doctor

■ do not take more than 6 caplets in 24 hours, unless directed by a doctor.
Questions or comments? 1-800-98-ADVIL.
Pfizer, Madison, NJ 07940 USA © 2020 Pfizer Inc.
For most recent product information, visit www.Advil.com

PAA118120.FDA02

UNVARNISHED AND PRINTED IN BLACK OR DARK BLUE FOR LOT & EXPIRATION AREA.

PAA118120.FDA02

MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.95 pt
• BULLETS	3.5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	.5 pt

Drug Facts (continued) ■ if you have ever had an allergic reaction to acetaminophen or any other pain reliever ■ right before or after heart surgery Ask a doctor before use if ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic	Drug Facts (continued) Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug When using this product ■ take with food or milk if stomach upset occurs Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
---	--

PAIN17961, FDA/02

LIFT HERE

<i>Drug Facts</i> (continued) ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ redness or swelling is present in the painful area ■ any new symptoms appear If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.	<i>Drug Facts</i> (continued) Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. <i>Directions</i> ■ do not take more than directed <table border="1"> <tr> <td>adults and children 12 years and over</td> <td>■ take 2 caplets every 8 hours while symptoms persist</td> </tr> <tr> <td>children under 12 year</td> <td>■ ask a doctor</td> </tr> </table> ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor <i>Other information</i> ■ read all warnings and directions before use. Keep carton.	adults and children 12 years and over	■ take 2 caplets every 8 hours while symptoms persist	children under 12 year	■ ask a doctor
adults and children 12 years and over	■ take 2 caplets every 8 hours while symptoms persist				
children under 12 year	■ ask a doctor				

<i>Drug Facts</i> (continued)
■ store at 20-25°C (68-77°F)
■ avoid excessive heat above 40°C (104°F)
<i>Inactive ingredients</i> carnauba wax, colloidal silicon dioxide, croscarmellose sodium, ferric oxides, glyceryl dibehenate, hypromellose, pharmaceutical ink, polydextrose, polyethylene glycol, pregelatinized starch, titanium dioxide
<i>Questions or comments?</i> call weekdays 9 AM to 5 PM at 1-800-88-ADVL

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	N/A
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

CONTAINS 2 MEDICINES

Advil

DUAL ACTION

WITH Acetaminophen

Acetaminophen 250 mg and Ibuprofen (NSAID) 125 mg Tablets Pain Reliever

Advil II

144 Caplets

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Uses

temporarily relieves minor aches and pains due to: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ muscular aches ■ minor pain of arthritis

Warnings

Acetaminophen liver damage warning:

This product contains acetaminophen. Severe liver damage may occur if you take with other drugs containing acetaminophen; more than 6 caplets in 24 hours, which is the maximum daily amount for this product; 3 or more alcoholic drinks every day while using this product.

Acetaminophen allergy alert:

may cause severe skin reactions. Symptoms may include: skin reddening, blisters, and rash. If skin reaction occurs, stop use and seek medical help right away.

NSAID allergy alert:

ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Do not use this product

if you have ever had an allergic reaction to any pain reliever; with any drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist; or right before or after surgery.

NSAID stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinning (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed.

Heart attack and stroke warning:

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Ask a doctor before use

if you are pregnant or breast feeding; you have liver disease; stomach bleeding warning applies to you; you have problems or serious side effects from taking pain relievers; you have a history of stomach problems, such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; you are taking a diuretic.

Ask your doctor or pharmacist before use

if you are under a doctor's care for any serious condition; taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin; or taking any other drug.

When using this product

take with food or milk if stomach upset occurs.

Stop use and ask a doctor

if pain gets worse or lasts more than 10 days; redness or swelling is present in the painful area; or any new symptoms appear.

Directions

■ do not take more than directed

adults and children 12 years and over

take 2 caplets every 8 hours while symptoms persist

children under 12 years

ask a doctor

■ do not take more than 6 caplets in 24 hours, unless directed by a doctor

Questions or comments?

1-800-88-ADVIL

Pfizer, Madison, NJ 07940 USA

© 2020 Pfizer Inc.

For most recent product information, visit www.Advil.com

PAA116862.FDA02

← UNVARNISHED AND PRINTED IN BLACK OR DARK BLUE FOR LOT & EXPIRATION AREA.

PAA116862.FDA02

MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	5 pt
• BULLETS	3.5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	.5 pt

TOP
PANEL 1

NEW

Easy Open ARTHRITIS CAP

CONTAINS 2 MEDICINES

Acetaminophen + Ibuprofen

Arthritis Foundation

Ease of Use™

CAP

Advil®

DUAL ACTION

WITH Acetaminophen

Acetaminophen 250 mg and
Ibuprofen (NSAID) 125 mg Tablets

Pain Reliever

Advil II

144 Caplets*

*Capsule-Shaped Tablets

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Do not use with other medicines containing ACETAMINOPHEN; can cause liver damage.

Drug Facts

Active ingredients (in each caplet) Purposes

Acetaminophen 250 mgPain reliever

Ibuprofen 125 mg (NSAID)*Pain reliever

*nonsteroidal anti-inflammatory drug

Uses

temporarily relieves minor aches and pains due to:

headache

toothache

backache

menstrual cramps

muscular aches

minor pain of arthritis

PAA116863.FDA02

81% For Position Only

3 0573 0147 897

LIFT HERE

For More Drug Facts

FOR LOT & EXPIRATION AREA.
UNVARNISHED AND PRINTED
IN BLACK OR DARK BLUE.

BASE
PANEL 3

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

under a doctor's care for any serious condition

taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin

taking any other drug

When using this product

take with food or milk if stomach upset occurs

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

feel faint

vomit blood

have bloody or black stools

have stomach pain that does not get better

you have symptoms of heart problems or stroke:

chest pain

trouble breathing

weakness in one part or side of body

slurred speech

leg swelling

pain gets worse or lasts more than 10 days

redness or swelling is present in the painful area

any new symptoms appear

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

do not take more than directed

adults and children 12 years and over

take 2 caplets every 8 hours while symptoms persist

children under 12 years

ask a doctor

do not take more than 6 caplets in 24 hours, unless directed by a doctor

Other information

read all warnings and directions before use.

store at 20-25°C (68-77°F)

avoid excessive heat above 40°C (104°F)

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, ferric oxides, glyceryl dibehenate, hypromellose, pharmaceutical ink, polydextrose, polyethylene glycol, pregelatinized starch, titanium dioxide

Questions or comments?

call weekdays 9 AM to 5 PM at 1-800-88-ADVIL

For most recent product information, visit [www.Advil.com](#)

Pfizer, Madison, NJ 07940 USA

For US Patent or Application status see [www.productpats.com](#)

© 2020 Pfizer Inc.

RELEASED BACK
PANEL 2

Drug Facts (continued)

Warnings

Acetaminophen liver damage warning:

This product contains acetaminophen. Severe liver damage may occur if you take:

with other drugs containing acetaminophen

more than 6 caplets in 24 hours, which is the maximum daily amount for this product

3 or more alcoholic drinks every day while using this product

Acetaminophen allergy alert:

may cause severe skin reactions. Symptoms may include:

skin reddening

blisters

rash

If skin reaction occurs, stop use and seek medical help right away.

NSAID allergy alert:

ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

hives

facial swelling

asthma (wheezing)

shock

skin reddening

rash

blisters

If an allergic reaction occurs, stop use and seek medical help right away.

NSAID stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

are age 60 or older

have had stomach ulcers or bleeding problems

take a blood thinning (anticoagulant) or steroid drug

Drug Facts (continued)

take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

have 3 or more alcoholic drinks every day while using this product

take more or for a longer time than directed

Heart attack and stroke warning:

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

with any other drug containing acetaminophen (prescription or nonprescription).

If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

if you have ever had an allergic reaction to acetaminophen or any other pain reliever

right before or after heart surgery

Ask a doctor before use if

you have liver disease

stomach bleeding warning applies to you

you have problems or serious side effects from taking pain relievers

you have a history of stomach problems, such as heartburn

you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

you are taking a diuretic

PAA116863.FDA02

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

090177e192dae21d\Final\Final On: 17-Feb-2020 15:53 (GMT)



DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

CONTAINS 2 MEDICINES

Advil

DUAL ACTION

WITH Acetaminophen

Acetaminophen 250 mg and
Ibuprofen (NSAID) 125 mg Tablets
Pain Reliever

Advil II

162
Caplets

READ AND KEEP CARTON FOR COMPLETE
WARNINGS AND INFORMATION

Do Not Use if seal under bottle cap imprinted with
"SEALED for YOUR PROTECTION" is broken or missing.

It may temporarily relieve minor aches and pains due to:

■ headache ■ toothache

■ backache ■ menstrual cramps ■ muscular aches ■ minor pain of arthritis

Warnings

Acetaminophen liver damage warning: This product contains acetaminophen. Severe liver damage may occur if you take: with other drugs containing acetaminophen; more than 6 caplets in 24 hours, which is the maximum daily amount for this product; 3 or more alcoholic drinks every day while using this product.

Acetaminophen allergy alert: may cause severe skin reactions. Symptoms may include: skin reddening, blisters, and rash. If skin reaction occurs, stop use and seek medical help right away.

NSAID allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Do not use this product if you have ever had an allergic reaction to any pain reliever; with any drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist; or right before or after surgery.

NSAID stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinning (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Ask a doctor before use if you are pregnant or breast feeding; you have liver disease; stomach bleeding warning applies to you; you have problems or serious side effects from taking pain relievers; you have a history of stomach problems, such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; you are taking a diuretic.

Ask your doctor or pharmacist before use if you are under a doctor's care for any serious condition; taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin; or taking any other drug.

When using this product take with food or milk if stomach upset occurs.

Stop use and ask a doctor if pain gets worse or lasts more than 10 days; redness or swelling is present in the painful area; or any new symptoms appear.

Directions

■ do not take more than directed

adults and children
12 years and over

take 2 caplets every 8 hours
while symptoms persist

children under 12 years

ask a doctor

■ do not take more than 6 caplets in 24 hours, unless directed by a doctor

Questions or comments? 1-800-88-ADVIL

Pfizer, Madison, NJ 07940 USA

© 2020 Pfizer Inc.

For most recent product information, visit www.Advil.com

PAA117785.FDA02

UNVARNISHED AND PRINTED
IN BLACK OR DARK BLUE FOR
LOT & EXPIRATION AREA.

PAA117785.FDA02

MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	5 pt
• BULLETS	3.5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	.5 pt

TOP
PANEL 1

18 FREE
CAPLETS

Easy Open
ARTHRITIS CAP

Arthritis Foundation
Ease of
Use™
CAP

CONTAINS 2 MEDICINES
Acetaminophen + Ibuprofen

Advil®
DUAL ACTION
WITH Acetaminophen

Advil II
Advil II

Acetaminophen 250 mg and
Ibuprofen (NSAID) 125 mg Tablets
Pain Reliever

144
162
Caplets*

*Capsule-Shaped Tablets

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

Do Not Use if seal under bottle cap imprinted with
“SEALED for YOUR PROTECTION” is broken or missing.

Do not use with other medicines containing
ACETAMINOPHEN; can cause liver damage.

Drug Facts

Active ingredients (in each caplet) Purposes

Acetaminophen 250 mgPain reliever

Ibuprofen 125 mg (NSAID*)Pain reliever

*nonsteroidal anti-inflammatory drug

Uses

temporarily relieves minor aches and pains due to:

headache

toothache

backache

menstrual cramps

muscular aches

minor pain of arthritis

PAA116864.FDA02

81% For Position Only

3 0573 0147 880

LIFT HERE
For More Drug Facts

FOR LOT & EXPIRATION AREA.
UNVARNISHED AND PRINTED
IN BLACK OR DARK BLUE.

BASE
PANEL 3

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

under a doctor's care for any serious condition

taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin

taking any other drug

When using this product

take with food or milk if stomach upset occurs

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

feel faint

vomit blood

have bloody or black stools

have stomach pain that does not get better

you have symptoms of heart problems or stroke:

chest pain

trouble breathing

weakness in one part or side of body

slurred speech

leg swelling

pain gets worse or lasts more than 10 days

redness or swelling is present in the painful area

any new symptoms appear

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

do not take more than directed

adults and children 12 years and over

take 2 caplets every 8 hours while symptoms persist

children under 12 years

ask a doctor

do not take more than 6 caplets in 24 hours, unless directed by a doctor

Other information

read all warnings and directions before use.

store at 20-25°C (68-77°F)

avoid excessive heat above 40°C (104°F)

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, ferric oxides, glyceryl dibehenate, hypromellose, pharmaceutical ink, polydextrose, polyethylene glycol, pregelatinized starch, titanium dioxide

Questions or comments?

call weekdays 9 AM to 5 PM at 1-800-88-ADVIL

For most recent product information, visit www.Advil.com

Pfizer, Madison, NJ 07940 USA

For US Patent or Application status see www.productpats.com

© 2020 Pfizer Inc.

RELEASED BACK
PANEL 2

Drug Facts (continued)

Warnings

Acetaminophen liver damage warning:

This product contains acetaminophen. Severe liver damage may occur if you take:

with other drugs containing acetaminophen

more than 6 caplets in 24 hours, which is the maximum daily amount for this product

3 or more alcoholic drinks every day while using this product

Acetaminophen allergy alert:

may cause severe skin reactions. Symptoms may include:

skin reddening

blisters

rash

If skin reaction occurs, stop use and seek medical help right away.

NSAID allergy alert:

ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

hives

facial swelling

asthma (wheezing)

shock

skin reddening

rash

blisters

If an allergic reaction occurs, stop use and seek medical help right away.

NSAID stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

are age 60 or older

have had stomach ulcers or bleeding problems

take a blood thinning (anticoagulant) or steroid drug

Drug Facts (continued)

take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

have 3 or more alcoholic drinks every day while using this product

take more or for a longer time than directed

Heart attack and stroke warning:

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

with any other drug containing acetaminophen (prescription or nonprescription).

If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

if you have ever had an allergic reaction to acetaminophen or any other pain reliever

right before or after heart surgery

Ask a doctor before use if

you have liver disease

stomach bleeding warning applies to you

you have problems or serious side effects from taking pain relievers

you have a history of stomach problems, such as heartburn


you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

you are taking a diuretic


PAA116864.FDA02

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces


TOP
PANEL 1



BASE
PANEL 3



RELEASED BACK
PANEL 2



PAA118115.FDA02

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

CONTAINS 2 MEDICINES

Advil®

DUAL ACTION

WITH Acetaminophen

Acetaminophen 250 mg and Ibuprofen (NSAID) 125 mg Tablets Pain Reliever

216 Caplets

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Uses temporarily relieves minor aches and pains due to: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ muscular aches ■ minor pain of arthritis

Warnings

Acetaminophen liver damage warning:

This product contains acetaminophen. Severe liver damage may occur if you take: with other drugs containing acetaminophen; more than 6 caplets in 24 hours, which is the maximum daily amount for this product; 3 or more alcoholic drinks every day while using this product.

Acetaminophen allergy alert: may cause severe skin reactions. Symptoms may include: skin reddening, blisters, and rash. If skin reaction occurs, stop use and seek medical help right away.

NSAID allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Do not use this product if you have ever had an allergic reaction to any pain reliever; with any drug containing acetaminophen (prescription or nonprescription); if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist; or right before or after surgery.

NSAID stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinning (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Ask a doctor before use if you are pregnant or breast feeding; you have liver disease; stomach bleeding warning applies to you; you have problems or serious side effects from taking pain relievers; you have a history of stomach problems, such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; you are taking a diuretic.

Ask your doctor or pharmacist before use if you are under a doctor's care for any serious condition; taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin; or taking any other drug.

When using this product take with food or milk if stomach upset occurs.

Stop use and ask a doctor if pain gets worse or lasts more than 10 days; redness or swelling is present in the painful area; or any new symptoms appear.

Directions

■ do not take more than directed

adults and children 12 years and over	■ take 2 caplets every 8 hours while symptoms persist
children under 12 years	■ ask a doctor

■ do not take more than 6 caplets in 24 hours, unless directed by a doctor

Questions or comments? 1-800-88-ADVIL

Pfizer, Madison, NJ 07940 USA © 2020 Pfizer Inc. For most recent product information, visit www.Advil.com


PAA116931.FDA02

UNVARNISHED AND PRINTED
IN BLACK OR DARK BLUE FOR
LOT & EXPIRATION AREA.


PAA116931.FDA02

NON MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	5 pt
• SUBHEADINGS/BODY TEXT	5 pt
• LEADING	5.5 pt
• BULLETS	4 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	.5 pt


TOP
PANEL 1



BASE
PANEL 3




RELEASED BACK
PANEL 2



PAA116865.FDA02

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



CONTAINS 2 MEDICINES

Advil[®]

DUAL ACTION

WITH Acetaminophen

Acetaminophen 250 mg and Ibuprofen (NSAID) 125 mg Tablets Pain Reliever

Advil II

240 Caplets

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Uses

temporarily relieves minor aches and pains due to: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ muscular aches ■ minor pain of arthritis

Warnings

Acetaminophen liver damage warning:

This product contains acetaminophen. Severe liver damage may occur if you take: with other drugs containing acetaminophen; more than 6 caplets in 24 hours, which is the maximum daily amount for this product; 3 or more alcoholic drinks every day while using this product.

Acetaminophen allergy alert:

may cause severe skin reactions. Symptoms may include: skin reddening, blisters, and rash. If skin reaction occurs, stop use and seek medical help right away.

NSAID allergy alert:

ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Do not use this product if you have ever had an allergic reaction to any pain reliever; with any drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist; or right before or after surgery.

NSAID stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinning (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]; have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed.

Heart attack and stroke warning:

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Ask a doctor before use if you are pregnant or breast feeding; you have liver disease; stomach bleeding warning applies to you; you have problems or serious side effects from taking pain relievers; you have a history of stomach problems, such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; you are taking a diuretic.

Ask your doctor or pharmacist before use if you are under a doctor's care for any serious condition; taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin; or taking any other drug.

When using this product take with food or milk if stomach upset occurs.

Stop use and ask a doctor if pain gets worse or lasts more than 10 days; redness or swelling is present in the painful area; or any new symptoms appear.

Directions

■ do not take more than directed

adults and children 12 years and over	■ take 2 caplets every 8 hours while symptoms persist
children under 12 years	■ ask a doctor

■ do not take more than 6 caplets in 24 hours, unless directed by a doctor

Questions or comments?
1-800-88-ADVIL
Pfizer, Madison, NJ 07940 USA
© 2020 Pfizer Inc.
For most recent product information, visit www.Advil.com

PAA116857.FDA02

UNVARNISHED AND PRINTED IN BLACK OR DARK BLUE. FOR LOT & EXPIRATION AREA

PAA116857.FDA02

NON MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	6 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	.5 pt

CONTAINS 2 MEDICINES
Acetaminophen + Ibuprofen

NEW

Advil®
DUAL ACTION
WITH Acetaminophen

Acetaminophen 250 mg and
Ibuprofen (NSAID) 125 mg Tablets
Pain Reliever

240
Caplets*
*Capsule-Shaped Tablets

Do Not Use if seal under bottle cap imprinted with
"SEALED for YOUR PROTECTION" is broken or missing.

Do not use with other medicines containing
ACETAMINOPHEN; can cause liver damage.

Drug Facts

Active ingredients (in each caplet)

Purposes

Acetaminophen 250 mg Pain reliever

Ibuprofen 125 mg (NSAID*) Pain reliever

*nonsteroidal anti-inflammatory drug

Uses

temporarily relieves minor aches and pains due to:

headache

toothache

backache

menstrual cramps

muscular aches

minor pain of arthritis

LIFT HERE
for More Drug Facts

PAA116858.FDA03

81% For Position Only

L3 0573 0147 972

UNVARNISHED &
PRINTED AREA
FOR LOT & EXP.
(LINES DO NOT PRINT)

Drug Facts (continued)

Warnings

Acetaminophen liver damage warning:

This product contains acetaminophen. Severe liver damage may occur if you take:
with other drugs containing acetaminophen
more than 6 caplets in 24 hours, which is the maximum daily amount for this product
3 or more alcoholic drinks every day while using this product

Acetaminophen allergy alert: may cause severe skin reactions. Symptoms may include:
skin reddening blisters rash
If skin reaction occurs, stop use and seek medical help right away.

NSAID allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
hives facial swelling asthma (wheezing)
shock skin reddening rash blisters
If an allergic reaction occurs, stop use and seek medical help right away.

NSAID stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
are age 60 or older
have had stomach ulcers or bleeding problems
take a blood thinning (anticoagulant) or steroid drug
take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
have 3 or more alcoholic drinks every day while using this product
take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
if you have ever had an allergic reaction to acetaminophen or any other pain reliever
right before or after heart surgery

Ask a doctor before use if

you have liver disease
stomach bleeding warning applies to you
you have problems or serious side effects from taking pain relievers
you have a history of stomach problems, such as heartburn
you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
you are taking a diuretic

Ask a doctor or pharmacist before use if you are

under a doctor's care for any serious condition
taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
taking any other drug

When using this product

take with food or milk if stomach upset occurs

Drug Facts (continued)

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:
feel faint
vomit blood
have bloody or black stools
have stomach pain that does not get better

you have symptoms of heart problems or stroke:
chest pain
trouble breathing
weakness in one part or side of body
slurred speech
leg swelling
pain gets worse or lasts more than 10 days
redness or swelling is present in the painful area
any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

do not take more than directed

adults and children 12 years and over	take 2 caplets every 8 hours while symptoms persist
children under 12 years	ask a doctor

do not take more than 6 caplets in 24 hours, unless directed by a doctor

Other information

read all warnings and directions before use.
store at 20-25°C (68-77°F)
avoid excessive heat above 40°C (104°F)

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, ferric oxides, glyceryl dibehenate, hypromellose, pharmaceutical ink, polydextrose, polyethylene glycol, pregelatinized starch, titanium dioxide

Questions or comments?

call weekdays 9 AM to 5 PM at 1-800-88-ADVIL

For most recent product information, visit www.Advil.com
For US Patent or Application status see www.productpats.com
Pfizer, Madison, NJ 07940 USA © 2020 Pfizer Inc.

PAA116858.FDA03

NON MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

CONTAINS 2 MEDICINES

Acetaminophen + Ibuprofen

NEW

Advil

DUAL ACTION

WITH Acetaminophen

Acetaminophen 250 mg and
Ibuprofen (NSAID) 125 mg Tablets

Pain Reliever

288

Caplets*

*Capsule-Shaped Tablets

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Do not use with other medicines containing ACETAMINOPHEN; can cause liver damage.

Drug Facts

Active ingredients (in each caplet)

Purposes

Acetaminophen 250 mg.....Pain reliever

Ibuprofen 125 mg (NSAID*).....Pain reliever

*nonsteroidal anti-inflammatory drug

Uses

temporarily relieves minor aches and pains due to:

headache

toothache

backache

menstrual cramps

muscular aches

minor pain of arthritis

PAA116855.FDA03

81% For Position Only

L3 0573 0147 927

LIFT HERE
for More Drug Facts

UNVARNISHED &
PRINTED AREA
FOR LOT & EXP.
(LINES DO NOT PRINT)

Drug Facts (continued)

Warnings

Acetaminophen liver damage warning:

This product contains acetaminophen. Severe liver damage may occur if you take:

- with other drugs containing acetaminophen
- more than 6 caplets in 24 hours, which is the maximum daily amount for this product
- 3 or more alcoholic drinks every day while using this product

Acetaminophen allergy alert: may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If skin reaction occurs, stop use and seek medical help right away.

NSAID allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

NSAID stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you have ever had an allergic reaction to acetaminophen or any other pain reliever
- right before or after heart surgery

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Drug Facts (continued)

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

do not take more than directed

adults and children 12 years and over

take 2 caplets every 8 hours while symptoms persist

children under 12 years

ask a doctor

do not take more than 6 caplets in 24 hours, unless directed by a doctor

Other information

read all warnings and directions before use.

store at 20-25°C (68-77°F)

avoid excessive heat above 40°C (104°F)

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, ferric oxides, glyceryl dibehenate, hypromellose, pharmaceutical ink, polydextrose, polyethylene glycol, pregelatinized starch, titanium dioxide

Questions or comments?

call weekdays 9 AM to 5 PM at 1-800-88-ADVIL

For most recent product information, visit www.Advil.com
For US Patent or Application status see www.productpats.com
Pfizer, Madison, NJ 07940 USA © 2020 Pfizer Inc.

PAA116855.FDA03

NON MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Advil Dual Action 18ct or Larger Save \$1.00 IRC D000063752

090177e18fef2e82\Final\Final

OUTSIDE

CONSUMER: LIMIT ONE COUPON PER PURCHASE of products and quantities stated. LIMIT OF 4 LIKE COUPONS in same shopping trip. Coupons not authorized if purchasing products for resale. Void if transferred, sold, auctioned, reproduced or altered from original. Any other use constitutes fraud. You may pay sales tax. Do not send to: Pfizer Consumer Healthcare. **DEALER:** Sending to Pfizer Consumer Healthcare 616, NCH Marketing Services, P.O. Box 880001, El Paso, TX 88588-0001 signifies compliance with the "Pfizer Consumer Healthcare Redemption Policy", copy available by writing to the above address. Cash value is 1/100th of 1 cent. ©2019 Pfizer Inc D000063752

MANUFACTURER'S COUPON EXPIRES X/XX/XX

SAVE \$1.00

On any Advil® Dual Action 18ct or larger

INSIDE

MANUFACTURER'S COUPON EXPIRES X/XX/XX



Advil Dual Action 36ct or Larger Save \$2.00 IRC D000063754

OUTSIDE

CONSUMER: LIMIT ONE COUPON PER PURCHASE of products and quantities stated. LIMIT OF 4 LIKE COUPONS in same shopping trip. Coupons not authorized if purchasing products for resale. Void if transferred, sold, auctioned, reproduced or altered from original. Any other use constitutes fraud. You may pay sales tax. Do not send to Pfizer Consumer Healthcare. **DEALER:** Sending to Pfizer Consumer Healthcare 616, NCH Marketing Services, P.O. Box 880001, El Paso, TX 88588-0001 signifies compliance with the "Pfizer Consumer Healthcare Redemption Policy", copy available by writing to the above address. Cash value is 1/100th of 1 cent. ©2019 Pfizer Inc D000063754

MANUFACTURER'S COUPON EXPIRES X/XX/XX
SAVE \$2.00
On any Advil® Dual Action 36ct or larger

INSIDE

MANUFACTURER'S COUPON EXPIRES X/XX/XX



090177e18fef34c1\Final\Final

Advil Dual Action 72ct or Larger Save \$2.00 IRC D000063756

OUTSIDE

CONSUMER: LIMIT ONE COUPON PER PURCHASE of products and quantities stated. LIMIT OF 4 LIKE COUPONS in same shopping trip. Coupons not authorized if purchasing products for resale. Void if transferred, sold, auctioned, reproduced or altered from original. Any other use constitutes fraud. You may pay sales tax. Do not send to Pfizer Consumer Healthcare. **DEALER:** Sending to Pfizer Consumer Healthcare 616, NCH Marketing Services, P.O. Box 880001, El Paso, TX 88588-0001 signifies compliance with the "Pfizer Consumer Healthcare Redemption Policy", copy available by writing to the above address. Cash value is 1/100th of 1 cent. ©2019 Pfizer Inc. D000063756

MANUFACTURER'S COUPON EXPIRES X/XX/XX
SAVE \$2.00
On any Advil® Dual Action 72ct or larger

INSIDE

MANUFACTURER'S COUPON EXPIRES X/XX/XX



Advil Dual Action 72ct or Larger Save \$3.00 IRC D000063755

OUTSIDE

CONSUMER: LIMIT ONE COUPON PER PURCHASE of products and quantities stated. LIMIT OF 4 LIKE COUPONS in same shopping trip. Coupons not authorized if purchasing products for resale. Void if transferred, sold, auctioned, reproduced or altered from original. Any other use constitutes fraud. You may pay sales tax. Do not send to Pfizer Consumer Healthcare. **DEALER:** Sending to Pfizer Consumer Healthcare 616, NCH Marketing Services, P.O. Box 880001, El Paso, TX 88588-0001 signifies compliance with the "Pfizer Consumer Healthcare Redemption Policy", copy available by writing to the above address. Cash value is 1/100th of 1 cent.
©2019 Pfizer Inc. D000063755

MANUFACTURER'S COUPON EXPIRES X/XX/XX

SAVE \$3.00

On any Advil® Dual Action 72ct or larger

INSIDE

MANUFACTURER'S COUPON EXPIRES X/XX/XX



Advil Dual Action EZ Open 144ct Save \$2.00 IRC D000063757

OUTSIDE

CONSUMER: LIMIT ONE COUPON PER PURCHASE of products and quantities stated. LIMIT OF 4 LIKE COUPONS in same shopping trip. Coupons not authorized if purchasing products for resale. Void if transferred, sold, auctioned, reproduced or altered from original. Any other use constitutes fraud. You may pay sales tax. Do not send to Pfizer Consumer Healthcare. **DEALER:** Sending to Pfizer Consumer Healthcare 616, NCH Marketing Services, P.O. Box 880001, El Paso, TX 88588-0001 signifies compliance with the "Pfizer Consumer Healthcare Redemption Policy", copy available by writing to the above address. Cash value is 1/100th of 1 cent. ©2019 Pfizer Inc D000063757

MANUFACTURER'S COUPON EXPIRES X/XX/XX
SAVE \$2.00
On any Advil® Dual Action EZ Open 144ct

INSIDE


MANUFACTURER'S COUPON EXPIRES X/XX/XX



D000063753.FDA02



FRONT




BACK


D000064024.FDA02



D000064025.FDA02



D000064026.FDA02



This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

KAREN M MAHONEY
02/28/2020 02:05:43 PM