



Drug Facts (continued)

Other information

- store between 20° to 25°C (68° to 77°F). Protect from freezing.
- keep this carton and the enclosed User Guide for important information
- do not use if tape printed with "sealed for your protection" on top and bottom carton flaps is not intact

Inactive ingredients
benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose

Questions or comments?
1-800-317-2165
(Mon–Fri, 9AM–5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer.
The Astepro mark is owned by Viatis.
Used under license.
© 2021 Bayer.
Dist. by:
Bayer HealthCare LLC
Whippany, NJ 07981
Patents: patents.livewell.bayer.com
Product of Germany.
000000

INDOOR & OUTDOOR
Allergy Relief

NDC 0280-0065-02

ASTEPRO[®] ALLERGY
First Antihistamine Nasal Spray FDA-Approved for Over-the-Counter Use

ASTEPRO[®] ALLERGY
Azela C . C
ANTIHISTAMINE NASAL SPRAY

**FULL
PRESCRIPTION
STRENGTH**

**UP TO
24 HOUR
RELIEF OF**

- ◀ Nasal congestion
- ◀ Runny nose
- ◀ Sneezing
- ◀ Itchy nose



120 METERED SPRAYS
0.78 fl oz (23 mL)

Alcohol-Free
Fragrance-Free

**NEW!
ANTIHISTAMINE
NASAL SPRAY**

**STERIOD
FREE**



THIS WAY TO RELIEF

Drug Facts

Active ingredient (in each spray)	Purpose
Azelastine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine)	Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose

Warnings
Only for use in the nose. Do not spray in eyes or mouth.
Do not use if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you
■ have had recent nose ulcers or nose surgery
■ have had a nose injury that has not healed

When using this product
■ drowsiness may occur
■ avoid alcoholic drinks
■ alcohol, sedatives, and tranquilizers may increase drowsiness
■ be careful when driving a motor vehicle or operating machinery
■ you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying.
■ nasal discomfort or sneezing may occur right after use
■ do not share this bottle with anyone else as this may spread germs

Stop use and ask a doctor if
■ an allergic reaction, such as a skin rash, to this product occurs
■ you have severe or frequent nosebleeds


If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
■ Read the User Guide for how to:
■ prime the bottle before first use
■ prime bottle again if not used for 3 or more days
■ use the spray
■ clean the spray nozzle if it gets clogged


adults and children 12 years and older	This product may be used either once or twice a day: ■ once daily: use 2 sprays in each nostril. OR ■ twice daily: use 1 or 2 sprays in each nostril every 12 hours ■ do not use more than 4 sprays in each nostril in a 24 hour period
children 6 years to 11 years	■ an adult should supervise use ■ 1 spray in each nostril every 12 hours ■ do not use more than 2 sprays in each nostril in a 24 hour period
children under 6 years	do not use




LOT:
EXP:YYYY.MM




















ASTEPRO[®]

ALLERGY

STERIOD
FREE

First Antihistamine Nasal Spray FDA-Approved for Over-the-Counter Use

Drug Facts	
Active ingredient (in each spray) Azelastine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine).....	Purpose Antihistamine
Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose	
Warnings Only for use in the nose. Do not spray in eyes or mouth. Do not use if you have ever had an allergic reaction to this product or any of its ingredients Ask a doctor before use if you ■ have had recent nose ulcers or nose surgery ■ have had a nose injury that has not healed When using this product ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying. ■ nasal discomfort or sneezing may occur right after use ■ do not share this bottle with anyone else as this may spread germs Stop use and ask a doctor if ■ an allergic reaction, such as a skin rash, to this product occurs ■ you have severe or frequent nosebleeds If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Drug Facts (continued)	
Directions ■ Read the User Guide for how to: ■ prime the bottle before first use ■ use the spray ■ prime bottle again if not used for 3 or more days ■ clean the spray nozzle if it gets clogged	
adults and children 12 years and older	This product may be used either once or twice a day: ■ once daily: use 2 sprays in each nostril; OR ■ twice daily: use 1 or 2 sprays in each nostril every 12 hours. ■ do not use more than 4 sprays in each nostril in a 24 hour period
children 6 years to 11 years	■ an adult should supervise use ■ 1 spray in each nostril every 12 hours ■ do not use more than 2 sprays in each nostril in a 24 hour period
children under 6 years	do not use
Other information ■ store between 20° to 25°C (68° to 77°F). Protect from freezing. ■ keep this label and the enclosed User Guide for important information ■ do not use if sealed package is torn or opened	
Inactive ingredients benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose	
Questions or comments? 1-800-317-2165 (Mon–Fri, 9AM–5PM EST)	

Bayer and the Bayer Cross are registered trademarks of Bayer.
The Astepro mark is owned by Viartis. Used under license.
© 2021 Bayer.
Dist. by:
Bayer HealthCare LLC
Whippany, NJ 07981
Patents: patents.livewell.bayer.com
Product of Germany.
000000

Bayer

LOT:
EXP: YYYY-MM

NDC 0280-0065-03





Actual Size 100%

FDA00342-1

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
06/17/2021 02:41:15 PM