

MEDICATION GUIDE

METHADONE HYDROCHLORIDE

Oral Solution USP

R_x only

Methadone Hydrochloride Oral Solution USP is:

- A strong prescription pain medicine that contains an opioid (narcotic) that is used to manage pain severe enough to require daily around-the-clock, long-term treatment with an opioid, when other pain treatments such as non-opioid pain medicines or immediate-release opioid medicines do not treat your pain well enough or you cannot tolerate them.
- A long-acting opioid pain medicine that can put you at risk for overdose and death. Even if you take your dose correctly as prescribed you are at risk for opioid addiction, abuse, and misuse than can lead to death.
- Not for use to treat pain that is not around-the-clock
- Also used to manage drug addiction.

Important information about Methadone:

- Get emergency help right away if you take too much methadone hydrochloride (overdose). When you first start taking methadone, when your dose is changed, or if you take too much (overdose), serious or life-threatening breathing problems that can lead to death may occur.
- Never give anyone your methadone. They could die from taking it. Store methadone away from children and in a safe place to prevent stealing or abuse. Selling or giving away methadone is against the law.

Do not take Methadone if you have:

- severe asthma, trouble breathing, or other lung problems.
- a bowel blockage or have narrowing of the stomach or intestines.

Before taking Methadone, tell your healthcare provider if you have a history of:

- head injury, seizures
- liver, kidney, thyroid problems
- problems urinating
- Pancreas or gallbladder problems
- abuse of street or prescription drugs, alcohol addiction, or mental health problems.

heart rhythm problems

(Long QT syndrome)

Tell your healthcare provider if you are:

- pregnant or planning to become pregnant. Prolonged use of methadone during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated.
- breastfeeding. Methadone passes into breast milk and may harm your baby.
- taking prescription or over-the-counter medicines, vitamins, or herbal supplements. Taking methadone with certain other medicines may cause serious side effects.

When taking Methadone:

- Do not change your dose. Take methadone exactly as prescribed by your healthcare provider.
- Do not take more than your prescribed dose in 24 hours. If you take methadone for pain and miss a dose, take methadone as soon as possible and then take your next dose 8 or 12 hours later as directed by your healthcare provider. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule.
- If you take methadone for opioid addiction and miss a dose, take your next dose the following day as scheduled. Do not take extra doses. Taking more than the prescribed dose may cause you to overdose because methadone builds up in your body over time.
- Do not crush, dissolve, snort or inject methadone because this may cause you to overdose or die.
- Call your healthcare provider if the dose you are taking does not control your pain.
- Do not stop taking methadone without talking to your healthcare provider.
- After you stop taking methadone, flush any unused tablets down the toilet.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Methadone Hydrochloride Oral Solution USP safely and effectively. See full prescribing information for Methadone Hydrochloride Oral Solution USP.

Methadone Hydrochloride Oral Solution USP, for oral use, CII
Initial U.S. Approval: 1947

WARNING: ADDICTION, ABUSE, AND MISUSE; LIFE-THREATENING RESPIRATORY DEPRESSION; ACCIDENTAL INGESTION; LIFE-THREATENING QT PROLONGATION; NEONATAL OPIOID WITHDRAWAL SYNDROME; AND TREATMENT FOR OPIOID ADDICTION
See full prescribing information for complete boxed warning

- Methadone Hydrochloride Oral Solution USP exposes users to risks of addiction, abuse, and misuse, which can lead to overdose and death. Assess each patient's risk before prescribing, and monitor regularly for development of these behaviors or conditions. (5.1, 9)
- Serious, life-threatening, or fatal respiratory depression may occur. Monitor closely, especially upon initiation or following a dose increase. (5.2)
- Accidental ingestion of Methadone Hydrochloride Oral Solution USP, especially in children, can result in fatal overdose of methadone. (5.2)
- QT interval prolongation and serious arrhythmia (torsades de pointes) have occurred during treatment with methadone. (5.3)
- Prolonged use of Methadone Hydrochloride Oral Solution USP during pregnancy can result in neonatal opioid withdrawal syndrome, which may be life-threatening if not recognized and treated. If opioid use is required for a prolonged period in a pregnant woman, advise the patient of the risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available (5.4).
- Methadone products, when used for the treatment of opioid addiction in detoxification or maintenance programs, shall be dispensed only by certified opioid treatment programs as stipulated in 42 CFR 8.12. (1)

RECENT MAJOR CHANGES
Boxed Warning 04/2014
Indications and Usage (1) 04/2014
Dosage and Administration (2) 04/2014
Warnings and Precautions (5) 04/2014

INDICATIONS AND USAGE

Methadone Hydrochloride Oral Solution USP is an opioid agonist indicated for the:

- Management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

- Limitations of Use**
- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with long-acting opioids, reserve Methadone Hydrochloride Oral Solution USP for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
 - Methadone Hydrochloride Oral Solution USP is not indicated as an as-needed (prn) analgesic.
 - Detoxification treatment of opioid addiction (heroin or other morphine-like drugs).
 - Maintenance treatment of opioid addiction (heroin or other morphine-like drugs), in conjunction with appropriate social and medical services. (1)

DOSAGE AND ADMINISTRATION
Management of Pain: For opioid-naïve patients, initiate methadone treatment with 2.5 mg every 8 to 12 hours. (2.2)

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: ADDICTION, ABUSE AND MISUSE; LIFE-THREATENING RESPIRATORY DEPRESSION; ACCIDENTAL INGESTION; LIFE-THREATENING QT PROLONGATION; NEONATAL OPIOID WITHDRAWAL SYNDROME; AND TREATMENT FOR OPIOID ADDICTION

- INDICATIONS AND USAGE
- DOSAGE AND ADMINISTRATION
 - Important General Information
 - Initial Dosing for Management of Pain
 - Titration and Maintenance of Therapy for Pain
 - Discontinuation of Methadone for Pain
 - Initiation/Initial Dosing for Detoxification and Maintenance Treatment of Opioid Addiction
 - Titration and Maintenance Treatment of Opioid Dependence Detoxification
 - Medically Supervised Withdrawal After a Period of Maintenance Treatment for Opioid Addiction
 - Risk of Relapse in Patients on Methadone Maintenance Treatment of Opioid Addiction
 - Considerations for Management of Acute Pain During Methadone Maintenance Treatment
- DOSAGE FORMS AND STRENGTHS
- CONTRAINDICATIONS
- WARNINGS AND PRECAUTIONS
 - Addiction, Abuse and Misuse
 - Life-Threatening Respiratory Depression
 - Life-Threatening QT Prolongation
 - Neonatal Opioid Withdrawal Syndrome
 - Interactions with Central Nervous System Depressants
 - Use in Elderly, Cachectic, and Debilitated Patients
 - Use in Patients with Chronic Pulmonary Disease
 - Hypotensive Effect
 - Use in Patients with Head Injury or Increased Intracranial Pressure
 - Use in Patients with Gastrointestinal Conditions
 - Use in Patients with Convulsive or Seizure Disorders

FULL PRESCRIBING INFORMATION:

WARNING: ADDICTION, ABUSE AND MISUSE; LIFE-THREATENING RESPIRATORY DEPRESSION; ACCIDENTAL INGESTION; LIFE-THREATENING QT PROLONGATION; NEONATAL OPIOID WITHDRAWAL SYNDROME; AND TREATMENT FOR OPIOID ADDICTION

Addiction, Abuse, and Misuse
Methadone Hydrochloride Oral Solution USP exposes patients and other users to the risks of opioid addiction, abuse, and misuse, which can lead to overdose and death. Assess each patient's risk prior to prescribing Methadone Hydrochloride Oral Solution USP, and monitor all patients regularly for the development of these behaviors or conditions [see *Warnings and Precautions* (5.1)].
Life-threatening Respiratory Depression
Serious, life-threatening, or fatal respiratory depression may occur with use of Methadone Hydrochloride Oral Solution USP. Monitor for respiratory depression, especially during initiation of Methadone Hydrochloride Oral Solution USP or following a dose

- To convert to methadone from another opioid, use available conversion factors to obtain estimated dose. (2.2)
- Initiation of Detoxification and Maintenance Treatment: A single dose of 20 to 30 mg may be sufficient to suppress withdrawal syndrome. (2.5)
- Do not abruptly discontinue methadone in a physically dependent patient. (2.4, 5.12)

DOSAGE FORMS AND STRENGTHS
Oral Solution: each 5 mL contains 5 mg or 10 mg of Methadone Hydrochloride Oral Solution USP. (3)

CONTRAINDICATIONS
Significant respiratory depression (4)
Acute or severe bronchial asthma (4)
Known or suspected paralytic ileus (4)
Hypersensitivity to methadone (4)

WARNINGS AND PRECAUTIONS
Respiratory Depression: The peak respiratory depressant effect typically occurs later, and persists longer than the peak analgesic effect. (5.2)
May cause QT interval prolongation and serious arrhythmia. (5.3)
Interactions with CNS depressants: Concomitant use may cause profound sedation, respiratory depression, and death. If coadministration is required, consider dose reduction of one or both drugs because of additive pharmacological effects. (5.5, 7.1)
Elderly, cachectic, debilitated patients and those with chronic pulmonary disease: Monitor closely because of increased risk for life-threatening respiratory depression. (5.6, 5.7)
Hypotensive effect: Monitor during dose initiation and titration. (5.8)
Patients with head injury or increased intracranial pressure: Monitor for sedation and respiratory depression. Avoid use of methadone in patients with impaired consciousness or coma susceptible to intracranial effects of CO₂ retention. (5.9)

ADVERSE REACTIONS
Most common adverse reactions are: lightheadedness, dizziness, sedation, nausea, vomiting, and sweating. (6)
To report SUSPECTED ADVERSE REACTIONS, contact Roxane Laboratories, Inc. at 1-800-952-8364 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS
CYP3A4 Inducers: Increased risk of more rapid metabolism and decreased effects of methadone. (7.2)
CYP3A4 Inhibitors: Increased risk of reduced metabolism and methadone toxicity. (7.2)
Anti-retroviral Agents: May result in increased clearance and decreased plasma levels of methadone or in certain cases, increased plasma levels and risk of toxicity. (7.2)
Potentially Arrhythmogenic Agents: Extreme caution is necessary when any drug known to have the potential to prolong the QT interval is prescribed in conjunction with methadone. (7.3)
Mixed Agonist/Antagonist and Partial Agonist Opioid Analgesics: Avoid use with methadone because they may reduce analgesic effect of methadone or precipitate withdrawal symptoms. (5.12, 7.4)

USE IN SPECIFIC POPULATIONS
Pregnancy: Based on animal data, may cause fetal harm. (8.1)
Nursing women: Methadone has been detected in human milk. Closely monitor infants of nursing women receiving methadone. (8.3)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide

April 2014

by neonatology experts. If opioid use is required for a prolonged period in a pregnant woman, advise the patient of the risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available [see *Warnings and Precautions* (5.4)].
Conditions For Distribution And Use Of Methadone Products For The Treatment Of Opioid Addiction
For detoxification and maintenance of opioid dependence, methadone should be administered in accordance with the treatment standards cited in 42 CFR Section 8, including limitations on unsupervised administration [see *Indications and Usage* (1)].

INDICATIONS AND USAGE

- Methadone Hydrochloride Oral Solution USP is indicated for the:
- Management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.
- Limitations of Use**
- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with long-acting opioids, reserve Methadone Hydrochloride Oral Solution USP for use in patients for whom alternative analgesic treatment options (e.g., non-opioid analgesics or immediate-release opioid analgesics) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
 - Methadone Hydrochloride Oral Solution USP is not indicated as an as-needed (prn) analgesic.
 - Detoxification treatment of opioid addiction (heroin or other morphine-like drugs).
 - Maintenance treatment of opioid addiction (heroin or other morphine-like drugs), in conjunction with appropriate social and medical services.

Conditions For Distribution And Use Of Methadone Products For The Treatment Of Opioid Addiction
Code of Federal Regulations, Title 42, Sec. 8
Methadone products when used for the treatment of opioid addiction in detoxification or maintenance programs, shall be dispensed only by opioid treatment programs (and agencies, practitioners or institutions by formal agreement with the program sponsor) certified by the Substance Abuse and Mental Health Services Administration and approved by the designated state authority. Certified treatment programs shall dispense and use methadone in oral form only and according to the treatment requirements stipulated in the Federal Opioid Treatment Standards (42 CFR 8.12). See below for important regulatory exceptions to the general requirement for certification to provide opioid agonist treatment.

Failure to abide by the requirements in these regulations may result in criminal prosecution, seizure of the drug supply, revocation of the program approval, and injunction precluding operation of the program.
Regulatory Exceptions To The General Requirement For Certification To Provide Opioid Agonist Treatment: During inpatient care, when the patient was admitted for any condition other than concurrent opioid addiction (pursuant to 21CFR 1306.07(c)), to facilitate the treatment of the primary admitting diagnosis). During an emergency period of no longer than 3 days while definitive care for the addiction is being sought in an appropriately licensed facility (pursuant to 21CFR 1306.07(b)).

DOSAGE AND ADMINISTRATION

Important General Information

- The peak respiratory depressant effect of methadone occurs later and persists longer than its peak therapeutic effect.
- A high degree of opioid tolerance does not eliminate the possibility of methadone overdose, iatrogenic or otherwise. Deaths have been reported during conversion to methadone from chronic, high-dose treatment with other opioid agonists and during initiation of methadone treatment of addiction in subjects previously abusing high doses of other agonists.
- With repeated dosing, methadone is retained in the liver and then slowly released, prolonging the duration of potential toxicity.
- Methadone has a narrow therapeutic index, especially when combined with other drugs.

Initial Dosing for Management of Pain

- Methadone should be prescribed only by healthcare professionals who are knowledgeable in the use of potent opioids for the management of chronic pain.
- Consider the following factors that differentiate methadone from other opioid analgesics:
 - There is high interpatient variability in absorption, metabolism, and relative analgesic potency. Population-based equianalgesic conversion ratios between methadone and other opioids are not accurate when applied to individuals.
 - The duration of analgesic action of methadone is 4 to 8 hours (based on single-dose studies) but the plasma elimination half-life is 8 to 59 hours.
 - Steady-state plasma concentrations, and full analgesic effects, are not attained until 3 to 5 days after initiation of dosing.

Initiate the dosing regimen for each patient individually, taking into account the patient's prior analgesic treatment experience and risk factors for addiction, abuse, and misuse [see *Warnings and Precautions* (5.1)]. Monitor patients closely for respiratory depression, especially within the first 24-72 hours of initiating therapy with methadone [see *Warnings and Precautions* (5.2)].
Use of Methadone as the First Opioid Analgesic: Initiate treatment with methadone with 2.5 mg orally every 8 to 12 hours.

Conversion from Other Oral Opioids to Methadone: Discontinue all other around-the-clock opioid drugs when methadone therapy is initiated. **Deaths have occurred in opioid-tolerant patients during conversion to methadone.**

While there are useful tables of opioid equivalents readily available, there is substantial inter-patient variability in the relative potency of different opioid drugs and products. As such, it is safer to underestimate a patient's 24-hour oral methadone requirements and provide rescue medication (e.g., immediate-release opioid) than to overestimate the 24-hour oral methadone requirements which could result in adverse reactions. With repeated dosing, the potency of methadone increases due to systemic accumulation.

- This is **not** a table of equianalgesic dose.
- The conversion factors in this table are only for the conversion **from** another oral opioid analgesic to methadone.
- The table **cannot** be used to convert **from** methadone **to** another opioid. Doing so will result in an overestimation of the dose of the new opioid and may result in fatal overdose.

Table 1: Conversion Factors to Methadone

Total Daily Baseline Oral Morphine Equivalent Dose	Estimated Daily Oral Methadone Requirement as Percent of Total Daily Morphine Equivalent Dose
< 100 mg	20% to 30%
100 to 300 mg	10% to 20%
300 to 600 mg	8% to 12%
600 mg to 1000 mg	5% to 10%
> 1000 mg	< 5 %

To calculate the estimated methadone dose using Table 1:

- For patients on a single opioid, sum the current total daily dose of the opioid, convert it to a Morphine Equivalent Dose according to specific conversion factor for that specific opioid, then multiply the Morphine Equivalent Dose by the corresponding percentage in the above table to calculate the approximate oral methadone daily dose. Divide the total daily methadone dose derived from the table above to reflect the intended dosing schedule (i.e., for administration every 8 hours, divide total daily methadone dose by 3).
- For patients on a regimen of more than one opioid, calculate the approximate oral methadone dose for each opioid and sum the totals to obtain the approximate total methadone daily dose. Divide the total daily methadone dose derived from the table above to reflect the intended dosing schedule (i.e., for administration every 8 hours, divide total daily methadone dose by 3).
- For patients on a regimen of fixed-ratio opioid/non-opioid analgesic products, use only the opioid component of these products in the conversion.
Always round the dose down, if necessary, to the appropriate methadone strength(s) available.
Example conversion from a single opioid to methadone:

- Step 1:** Sum the total daily dose of the opioid (in this case, Morphine Extended Release Tablets 50 mg twice daily)
50 mg Morphine Extended Release Tablets 2 times daily = 100 mg total daily dose of Morphine
- Step 2:** Calculate the approximate equivalent dose of Methadone Hydrochloride Oral Solution USP based on the total daily dose of Morphine using Table 1.
100 mg total daily dose of Morphine x 15% (10% to 20% per Table 1) = 15 mg Methadone Hydrochloride Oral Solution USP daily

Step 3: Calculate the approximate starting dose of Methadone Hydrochloride Oral Solution USP to be given every 12 hours. Round down, if necessary, to the appropriate methadone tablets strengths available.

15 mg daily / 2 = 7.5 mg Methadone Hydrochloride Oral Solution USP every 12 hours
Then 7.5 mg is rounded down to 5 mg Methadone Hydrochloride Oral Solution USP every 12 hours
Close observation and frequent titration are warranted until pain management is stable on the new opioid. Monitor patients for signs and symptoms of opioid withdrawal or for signs of over-sedation/toxicity after converting patients to Methadone Hydrochloride Oral Solution USP.
Conversion from Parenteral Methadone to Methadone Hydrochloride Oral Solution USP: Use a conversion ratio of 1:2 mg for parenteral to oral methadone (e.g., 5 mg parenteral methadone to 10 mg oral methadone).

Titration and Maintenance of Therapy for Pain

Individually titrate methadone to a dose that provides adequate analgesia and minimizes adverse reactions. Continually reevaluate patients receiving methadone to assess the maintenance of pain control and the relative incidence of adverse reactions, as well as monitoring for the development of addiction, abuse, or misuse. Frequent communication is important among the prescriber, other members of the healthcare team, the patient, and the caregiver to ensure that the patient's pain is managed with the first dose of analgesic. During chronic therapy, periodically reassess the continued need for the use of opioid analgesics.

Because steady-state plasma concentrations are approximated within 24 to 36 hours, methadone dosage adjustments may be done every 1 to 2 days.
Patients who experience breakthrough pain may require a dose increase of methadone, or may need rescue medication with an appropriate dose of an immediate-release medication. If the level of pain increases after dose stabilization, attempt to identify the source of increased pain before increasing the methadone dose.

If unacceptably opioid-related adverse reactions are observed, subsequent doses may be reduced and/or the dosing interval adjusted (i.e., every 8 hours or every 12 hours). Adjust the dose to obtain an appropriate balance between management of pain and opioid-related adverse reactions.

Discontinuation of Methadone for Pain

When a patient no longer requires therapy with methadone for pain, use a gradual downward titration, of the dose every two to four days, to prevent signs and symptoms of withdrawal in the physically-dependent patient. Do not abruptly discontinue methadone.

Induction/Initial Dosing for Detoxification and Maintenance Treatment of Opioid Addiction

For detoxification and maintenance of opioid dependence methadone should be administered in accordance with the treatment standards cited in 42 CFR Section 8.12, including limitations on unsupervised administration.

Administer the initial methadone dose under supervision, when there are no signs of sedation or intoxication, and the patient shows symptoms of withdrawal. An initial single dose of 20 to 30 mg of methadone will often be sufficient to suppress withdrawal symptoms. The initial dose should not exceed 30 mg.

To make same-day dosing adjustments, have the patient wait 2 to 4 hours for further evaluation, when peak levels have been reached. Provide an additional 5 to 10 mg of methadone if withdrawal symptoms have not been suppressed or if symptoms reappear.

The total daily dose of methadone on the first day of treatment should not ordinarily exceed 40 mg. Adjust the dose over the first week of treatment based on control of withdrawal symptoms at the time of expected peak activity (e.g., 2 to 4 hours after dosing). When adjusting the dose, keep in mind that methadone levels will accumulate over the first several days of dosing; deaths have occurred in early treatment due to the cumulative effects. Instruct patients that the dose will "hold" for a longer period of time as tissue stores of methadone accumulate.

Use lower initial doses for patients whose tolerance is expected to be low at treatment entry. Any patient who has not taken opioids for more than 5 days may no longer be tolerant. Do not determine initial doses based on previous treatment episodes or dollars spent per day on illicit drug use.

Short-term treatment: For a brief course of stabilization followed by a period of medically supervised withdrawal, titrate the patient to a total daily dose of about 40 mg in divided doses to achieve an adequate stabilizing level. After 2 to 3 days of stabilization, gradually decrease the dose of methadone. Decrease the dose of methadone on a daily basis or at 2-day intervals, keeping the amount of methadone sufficient to keep withdrawal symptoms at a tolerable level. Hospitalized patients may tolerate a daily reduction of 20% of the total daily dose. Ambulatory patients may need a slower schedule.

Titration and Maintenance Treatment of Opioid Dependence Detoxification

Titrate patients in maintenance treatment to a dose that prevents opioid withdrawal symptoms for 24 hours, reduces drug hunger or craving, and blocks or attenuates the euphoric effects of self-administered opioids, ensuring that the patient is tolerant to the sedative effects of methadone. Most commonly, clinical stability is achieved at doses between 80 to 120 mg/day.

Medically Supervised Withdrawal After a Period of Maintenance Treatment for Opioid Addiction

There is considerable variability in the appropriate rate of methadone taper in patients choosing medically supervised withdrawal from methadone treatment. Dose reductions should generally be less than 10% of the established tolerance or maintenance dose, and 10 to 14-day intervals should elapse between dose reductions. Apprise patients of the risk of relapse to illicit drug use associated with discontinuation of methadone maintenance treatment.

Risk of Relapse in Patients on Methadone Maintenance Treatment of Opioid Addiction

Abrupt opioid discontinuation can lead to development of opioid withdrawal symptoms [see *Drug Abuse and Dependence* (9.3)]. Opioid withdrawal symptoms have been associated with an increased risk of relapse to illicit drug use in susceptible patients.

Considerations for Management of Acute Pain During Methadone Maintenance Treatment

Patients in methadone maintenance treatment for opioid dependence who experience physical trauma, post-operative pain or other acute pain cannot be expected to derive analgesia from their existing dose of methadone. Rescue pain should be administered analgesics, including opioids, in doses that would otherwise be indicated for non-methadone-treated patients with similar painful conditions. When opioids are required for management of acute pain in methadone maintenance patients, somewhat higher and/or more frequent doses will often be required than would be the case for non-tolerant patients due to the opioid tolerance induced by methadone.

Dosage Adjustment During Pregnancy

Methadone clearance may be increased during pregnancy. During pregnancy, a woman's methadone dose may need to be increased or the dosing interval decreased. Methadone should be used in pregnancy only if the potential benefit justifies the potential risk to the fetus [see *Use in Specific Populations* (8.1)].

DOSAGE FORMS AND STRENGTHS

Each 5 mL of ORANGE Methadone Hydrochloride Oral Solution USP contains methadone hydrochloride 5 mg or 10 mg. The concentration of the 5 mg per 5 mL solution is 1 mg/mL and the concentration of the 10 mg per 5 mL solution is 2 mg/mL.

CONTRAINDICATIONS

- Methadone Hydrochloride Oral Solution USP is contraindicated in patients with:
- Significant respiratory depression
 - Acute or severe bronchial asthma in an unmonitored setting or in the absence of resuscitative equipment
 - Known or suspected paralytic ileus
 - Hypersensitivity (e.g., anaphylaxis) to methadone [see *Adverse Reactions* (6)].

WARNINGS AND PRECAUTIONS

5.1 Addiction, Abuse and Misuse
Methadone Hydrochloride Oral Solution USP contains methadone, a Schedule II controlled substance. As an opioid, methadone exposes users to the risks of addiction, abuse, and misuse [see *Drug Abuse and Dependence* (9)]. As long-acting opioids such as methadone have pharmacological effects over an extended period of time, there is a greater risk for overdose and death.

Although the risk of addiction in any individual is unknown, it can occur in patients appropriately prescribed methadone and in those who obtain the drug illicitly. Addiction can occur at recommended doses and if the drug is misused or abused.

Assess each patient's risk for opioid addiction, abuse, or misuse prior to prescribing methadone, and monitor all patients receiving methadone for the development of these behaviors or conditions. Risks are increased in patients with a personal or family history of substance abuse (including drug or alcohol addiction or abuse) or mental illness (e.g., major depression). The potential for these risks should not, however, prevent the prescribing of methadone for the proper management of pain in any given patient. Patients at increased risk may be prescribed long-acting opioids such as methadone, but use in such patients necessitates intensive counseling about the risks and proper use of methadone along with the intensive monitoring for signs of addiction, abuse, and misuse.

Abuse or misuse of methadone by crushing, chewing, snorting, or injecting the dissolved product will result in the uncontrolled delivery of the methadone and can result in overdose and death [see *Overdose* (10)].

Opioid agonists such as methadone are sought by drug abusers and people with addiction disorders and are subject to criminal diversion. Consider these risks when prescribing or dispensing methadone. Strategies to reduce these risks include prescribing the drug in the smallest appropriate quantity and advising the patient on the proper disposal of unused drug [see *Patient Counseling Information* (17)]. Contact local state professional licensing board or state controlled substances authority for information on how to prevent and detect abuse or diversion of this product.

While taking Methadone DO NOT:

- Drive or operate heavy machinery, until you know how methadone affects you. Methadone can make you sleepy, dizzy, or lightheaded.
- Drink alcohol or use prescription or over-the-counter medicines that contain alcohol. Using products containing alcohol during treatment with methadone may cause you to overdose and die.

The possible side effects of Methadone are:

- constipation, nausea, sleepiness, vomiting, tiredness, headache, dizziness, abdominal pain. Call your healthcare provider if you have any of these symptoms and they are severe.

Get emergency medical help if you have:

- trouble breathing, shortness of breath, fast heartbeat, chest pain, swelling of your face, tongue or throat, extreme drowsiness, lightheadedness when changing positions, or you are feeling faint.

These are not all the possible side effects of methadone. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. **For more information go to dailymed.nlm.nih.gov.**

Roxane Laboratories, Inc. Columbus Ohio 43216, www.roxane.com or call 1-800-962-8364.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Roxane Laboratories, Inc.
Columbus, Ohio 43216

4056301/06

Revised April 2014
© RLI, 2014



The major hazards of methadone are respiratory depression, and to a lesser degree, systemic hypotension. Respiratory arrest, shock, cardiac arrest, and death have occurred.

The most frequently observed adverse reactions include lightheadedness, dizziness, sedation, nausea, vomiting and sweating. These effects seem to be more prominent in ambulatory patients and in those who are not suffering severe pain. In such individuals, lower doses are advisable.

Other adverse reactions include the following:

Body as a Whole: asthenia (weakness), edema, headache

Cardiovascular: arrhythmias, bigeminal rhythms, bradycardia, cardiomyopathy, ECG abnormalities, extrasystoles, flushing, heart failure, hypotension, palpitations, phlebitis, QT interval prolongation, syncope, T-wave inversion, tachycardia, torsades de pointes, ventricular fibrillation, ventricular tachycardia

Central Nervous System: agitation, confusion, disorientation, dysphoria, euphoria, insomnia, hallucinations, seizures, visual disturbances

Endocrine: hypogonadism

Gastrointestinal: abdominal pain, anorexia, biliary tract spasm, constipation, dry mouth, glossitis

Hematologic: reversible thrombocytopenia has been described in opioid addicts with chronic hepatitis

Metabolic: hypokalemia, hypomagnesemia, hypomagnesemia, muscle gain

Renal: antidiuretic effect, urinary retention or hesitancy

Reproductive: amenorrhea, reduced libido and/or potency, reduced ejaculate volume, reduced seminal vesicle and prostate secretions, decreased sperm motility, abnormalities in sperm morphology

Respiratory: pulmonary edema, respiratory depression

Skin and Subcutaneous Tissue: pruritus, urticaria, other skin rashes, and rarely, hemorrhagic urticaria

Hypersensitivity: Anaphylaxis has been reported with ingredients contained in methadone. Advise patients how to react to such attacks and what to seek medical attention.

Maintenance on a Stabilized Dose: During prolonged administration of methadone, as in a methadone maintenance treatment program, constipation and sweating often persist and hypogonadism, decreased serum testosterone and reproductive effects are thought to be related to chronic opioid use.

Methadone the Detoxification and Maintenance Treatment of Opioid Dependence: During the induction phase of methadone maintenance treatment, patients are being withdrawn from illicit opioids and may have opioid withdrawal symptoms. Monitor patients for signs and symptoms including: lacrimation, rhinorrhea, sneezing, yawning, excessive perspiration, goose-flesh, fever, chills alternating with flushing, restlessness, irritability, weak neonatal mortality and mortality when compared to controls. The dose of methadone may vary from a few days to weeks or even months. There is no consensus on the appropriate management of infant withdrawal [see *Dosage and Administration* (2.10)].

Effects on the Neonate: Babies born to mothers who have been taking opioids regularly prior to delivery may be physically dependent. Onset of withdrawal symptoms in infants is usually in the first days after birth. Monitor newborn for withdrawal signs and symptoms including: poor feeding, irritability, excessive crying, tremors, rigidity, hyper-areflexes, increased respiratory rate, diarrhea, sneezing, yawning, vomiting, fever, and seizures. The intensity of the neonatal withdrawal signs does not always correlate with the maternal dose or the duration of maternal exposure. The duration of the withdrawal signs may vary from a few days to weeks or even months. There is no consensus on the appropriate management of infant withdrawal [see *Warnings and Precautions* (5.4)].

Human Data: Reported studies have generally compared the benefit of methadone to the risk of untreated addiction to illicit drugs; the relevance of these findings to pain patients prescribed methadone during pregnancy is unclear. Pregnant women involved in methadone maintenance programs have been reported to have significantly improved prenatal care leading to significantly reduced incidence of obstetric complications, weaker neonatal mortality and mortality when compared to controls. Several factors, including maternal use of illicit drugs, nutrition, infection and psychosocial circumstances, complicate the interpretation of investigations of the children of women who take methadone during pregnancy. Information is limited regarding dose and duration of methadone use during pregnancy, and most maternal exposure appears to occur after the first trimester of pregnancy.

A review of published data on experiences with methadone use during pregnancy by the Teratogen Information System (TERIS) concluded that maternal use of methadone during pregnancy as part of a supervised, carefully monitored program poses a substantial, though small, risk (quantity and quality of data assessed as "limited to fair"). However, the data are insufficient to state there is no risk (TERIS, last reviewed October, 2002). A retrospective case series of 101 pregnant, opioid-dependent women who underwent inpatient opioid detoxification with methadone did not demonstrate any increased risk of miscarriage in the 2nd trimester or premature delivery in the 3rd trimester. Recent studies suggest an increased risk of premature delivery in opioid-dependent women exposed to methadone during pregnancy, although the presence of confounding factors makes it difficult to determine a causal relationship. Several studies have suggested that infants born to narcotic-addicted women treated with methadone during all or part of pregnancy had a higher risk of being born with a low birth weight, low Apgar scores, and/or head circumference compared to controls. This growth deficit does not appear to persist into later childhood. Children prenatally exposed to methadone have been reported to demonstrate mild but persistent deficits in performance on psychometric and behavioral tests. In addition, several studies suggest that children born to opioid-dependent women exposed to methadone during pregnancy may have an increased risk of visual development anomalies; however, a causal relationship has not been assigned.

There are conflicting reports on whether Sudden Infant Death Syndrome occurs with an increased incidence in infants born to women treated with methadone during pregnancy. Abnormal fetal non-stress tests were performed more frequently when the test is performed 1 to 2 hours after a maintenance dose of methadone in late pregnancy compared to controls.

Animal Data: Methadone did not produce teratogenic effects in rat or rabbit models. Methadone produced teratogenic effects following large doses, in the guinea pig, hamster and mouse. One published study in pregnant hamsters indicated that a single subcutaneous dose of methadone ranging from 31 to 185 mg/kg (the 31 mg/kg dose is approximately 2 times a human daily oral dose of 120 mg/day on a mg/m² basis) on day 8 of gestation resulted in a decrease in the number of fetuses per litter and an increase in the number of fetuses exhibiting congenital malformations described in the following order: craniofacial, cranial, and "various other lesions." The majority of the doses tested also resulted in maternal death. In another study, a single subcutaneous dose of 22 to 24 mg/kg methadone (estimated exposure was approximately equivalent to a human daily oral dose of 120 mg/day on a mg/m² basis) administered on day 9 of gestation in mice also produced exencephaly in 11% of the embryos. However, no effects were reported in rats and rabbits at oral doses up to 40 mg/kg (estimated exposure was approximately 3 and 6 times, respectively, a human daily oral dose of 120 mg/day on a mg/m² basis) administered during days 6 to 15 and 6 to 18, respectively.

Published animal data have reported increased neonatal mortality in the offspring of male rodents that were treated with methadone prior to mating. In these studies, the female rodents were not treated with methadone, indicating paternally-mediated developmental toxicity. Specifically, methadone administered to the male rat prior to mating with methadone-naïve females resulted in decreased weight gain in progeny after weaning. The male progeny demonstrated reduced thymus weights, whereas the female progeny demonstrated increased adrenal weights. Behavioral testing of these male and female progeny revealed significant differences in behavioral tests compared to control animals, suggesting that paternal methadone exposure produced physiological and behavioral changes in progeny in this model. Other animal studies have reported that perinatal exposure to opioids including methadone alters neuronal development and behavior in the offspring. Perinatal methadone exposure in rats has been linked to alterations in learning ability, motor activity, thermal regulation, nociceptive responses and sensitivity to drugs.

Additional animal data demonstrates evidence for neurochemical changes in the brains of methadone-treated offspring, including changes to the cholinergic, dopaminergic, noradrenergic and serotonergic systems. Studies demonstrated that methadone treatment of male rats for 21 to 32 days prior to mating with methadone-naïve females did not produce any adverse effects, suggesting that prolonged methadone treatment did not alter the response to the developmental toxicities noted in the progeny. Mechanistic studies in this rat model suggest that the developmental effects of "paternal" methadone on the progeny appear to be due to decreased testosterone production. These animal data mirror the reported clinical findings of decreased testosterone levels in human males on methadone maintenance therapy for opioid addiction and in males receiving chronic intrasplenic opioids.

Additional data have been published indicating that methadone treatment of male rats (once a day for three consecutive days) increased embryolethality and neonatal mortality. Examination of uterine contents of methadone-treated female mice bred to methadone-treated mice indicated that methadone treatment produced an increase in the rate of preimplantation deaths in all post-meiotic states.

8.2 Labor and Delivery

Opioids cross the placenta and may produce respiratory depression in neonates. Methadone is not for use in women during and immediately prior to labor, when shorter acting analgesics or other analgesic techniques are more appropriate. Opioid analgesics can prolong labor through actions that temporarily reduce the strength, duration, and frequency of uterine contractions. However this effect is not consistent and may be offset by an increased rate of cervical dilation, which tends to shorten labor.

8.3 Nursing Mothers

Methadone is secreted into human milk. At maternal oral doses of 10 to 80 mg/day, methadone concentrations from 50 to 570 mcg/L in milk have been reported, which, in the majority of samples, were lower than maternal serum drug concentrations at steady state. Peak methadone levels in milk occur approximately 4 to 5 hours after an oral dose. Based on an average milk consumption of 150 mL/kg/day, an infant would consume approximately 17.4 mcg/kg/day which is approximately 2 to 3% of the oral maternal dose. Methadone has been detected in very low plasma concentrations in some infants whose mothers were taking methadone. Cases of sedation and respiratory depression in infants exposed to methadone through breast milk have been reported. Caution should be exercised when methadone is administered to a nursing woman.

Advise women who are being treated with methadone and who are breastfeeding or express a desire to breastfeed of the presence of methadone in human milk. Instruct breastfeeding mothers how to identify respiratory depression and sedation in their babies and when it may be necessary to contact their healthcare provider or seek immediate medical care. Breastfed infants of mothers using methadone should be weaned gradually to prevent development of withdrawal symptoms in the infant.

8.4 Pediatric Use

The safety, effectiveness, and pharmacokinetics of methadone in pediatric patients below the age of 18 years have not been established.

8.5 Geriatric Use

Clinical studies of methadone did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently compared to younger subjects. Other reported clinical experience has not identified differences in responses between elderly and younger patients. In general, start elderly patients at the lower end of the dosing range, taking into account the greater frequency of decreased hepatic, renal, or cardiac function and of concomitant disease or other drug therapy in geriatric patients. Closely monitor elderly patients for signs of respiratory and central nervous system depression.

8.6 Renal Impairment

Methadone pharmacokinetics have not been extensively evaluated in patients with renal insufficiency. Since unmetabolized methadone and its metabolites are excreted in urine to a variable degree, start these patients on lower doses and with longer dosing intervals and titrate slowly while carefully monitoring for signs of respiratory and central nervous system depression.

8.7 Hepatic Impairment

Methadone has not been extensively evaluated in patients with hepatic insufficiency. Methadone is metabolized by hepatic pathways; therefore, patients with liver impairment may be at risk of increased systemic exposure to methadone after multiple dosing. Start these patients on lower doses and titrate slowly while carefully monitoring for signs of respiratory and central nervous system depression.

9 DRUG ABUSE AND DEPENDENCE

9.1 Controlled Substance

Methadone is a mu-agonist opioid with an abuse liability similar to other opioid agonists and is a Schedule II controlled substance. Methadone can be abused and is subject to misuse, addiction, and criminal diversion [see *Warnings and Precautions* (5.1)].

9.2 Abuse

All patients treated with opioids for pain management require careful monitoring for signs of abuse and addiction, since use of opioid analgesic products carries the risk of addiction even under appropriate medical use.

Abuse and addiction are separate and distinct from physical dependence and tolerance. Physicians should be aware that addiction may not be accompanied by concurrent tolerance and symptoms of physical dependence in all addicts. In addition, abuse of opioids can occur in the absence of true addiction.

Methadone, like other opioids, can be diverted for non-medical use into illicit channels of distribution. Careful record-keeping of prescribing information, including quantity, frequency, and renewal requests as required by state law, is strongly advised.

Risks Specific to Abuse of Methadone: Abuse of methadone poses a risk of overdose and death. This risk is increased with concurrent abuse of methadone and alcohol or other substances. Methadone is for oral use only and must not be injected. Parenteral drug abuse is commonly associated with transmission of infectious diseases such as hepatitis and HIV.

Proper assessment and selection of the patient, proper prescribing practices, periodic re-evaluation of therapy, and proper dispensing and storage are appropriate measures that help to limit abuse of opioid drugs.

9.3 Dependence

Both tolerance and physical dependence can develop during chronic opioid therapy. Tolerance is the need for increased doses of opioids to maintain a degree of analgesia (in the absence of disease progression or other external factors). Tolerance may occur to both the desired and undesired effects of drugs, and may develop at different rates for different effects.

Physical dependence results in withdrawal symptoms after abrupt discontinuation or a significant dose reduction of a drug. Withdrawal also may be precipitated during the administration of drugs with opioid antagonist activity, e.g., naloxone, mixed agonist/antagonist analgesics (pentazocine, butorphanol, nalbuphine), or partial agonists (buprenorphine). Physical dependence may not occur to a clinically significant degree until after several days to weeks of continued opioid usage.

Abuse and addiction are separate and distinct from physical dependence and tolerance. Physicians should be aware that addiction may not be accompanied by concurrent tolerance and symptoms of physical dependence in all addicts. In addition, abuse of opioids can occur in the absence of true addiction.

Methadone is abruptly discontinued in a physically dependent patient, an abstinence syndrome may occur. Some or all of the following can characterize this syndrome: restlessness, lacrimation, rhinorrhea, yawning, perspiration, chills, myalgia, and mydriasis. Other signs and symptoms also may develop, including irritability, anxiety, backache, joint pain, weakness, abdominal cramps, insomnia, nausea, anorexia, vomiting, diarrhea, or increased blood pressure, respiratory rate, or heart rate.

Infants born to mothers physically dependent on opioids will also be physically dependent and may exhibit respiratory difficulties and withdrawal symptoms [see *Use in Specific Populations* (8.1) and *Warnings and Precautions* (5.4)].

10 OVERDOSAGE

Clinical Presentation

Acute overdosage of methadone is manifested by respiratory depression, somnolence progressing to stupor or coma, maximally constricted pupils, skeletal-muscle flaccidity, cold and clammy skin, and, sometimes, bradycardia and hypotension. In severe overdosage, particularly by the intravenous route, apnea, circulatory collapse, cardiac arrest, and death may occur.

Treatment of Overdose

In case of overdose, priorities are the re-establishment of a patent and protected airway and institution of assisted or controlled ventilation if needed. Employ other supportive measures (including oxygen, vasopressors) in the management of circulatory shock and pulmonary edema as indicated. Cardiac arrest or arrhythmias will require advanced life support techniques.

The opioid antagonists, such as naloxone, are specific antidotes to respiratory depression resulting from opioid overdose. Opioid antagonists should not be administered in the absence of clinically significant respiratory or circulatory depression secondary to methadone overdose. Such agents should be administered cautiously to patients who are known, or suspected to be, physically dependent on methadone. In such cases, an abrupt or complete reversal of opioid effects may precipitate an acute withdrawal syndrome.

Because the duration of reversal would be expected to be less than the duration of action of methadone, carefully monitor the patient until spontaneous respiration is reliably re-established. If the response to opioid antagonists is suboptimal or not sustained, additional antagonist should be given as directed in the product's prescribing information.

Methadone, like other physically dependent on opioids, administration of an opioid receptor antagonist may precipitate an acute withdrawal. The severity of the withdrawal produced will depend on the degree of physical dependence and the dose of the antagonist administered. If a decision is made to treat serious respiratory depression in the physically dependent patient, administration of the antagonist should be begun with care and by titration with smaller than usual doses of the antagonist.

11 DESCRIPTION

Methadone hydrochloride is chemically described as 6-(dimethylamino)-4,4-diphenyl-3-heptanone hydrochloride. Methadone hydrochloride is a white, crystalline material that is water-soluble. Its molecular formula is C₂₁H₂₇NO•HCl and it has a molecular weight of 345.91. Methadone hydrochloride has a melting point of 235°C, and a pKa of 8.25 in water at 20°C. Its octanol/water partition coefficient at pH 7.4 is 117. A solution (1:100) in water has a pH between 4.5 and 6.5.

It has the following structural formula:

Each 5 mL of oral solution contains 5 or 10 mg of Methadone Hydrochloride USP and the following inactive ingredients: alcohol (8%), benzoic acid, citric acid, FD&C Red #40, FD&C Yellow #6, flavoring (lemon), glycerin, sorbitol, and water.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Methadone hydrochloride is a mu-agonist; a synthetic opioid analgesic with multiple actions qualitatively similar to those of morphine, the most prominent of which involves the central nervous system and organs composed of smooth muscle. The principal therapeutic uses for methadone are for analgesia and for the treatment of pain or the relief of withdrawal symptoms. The methadone withdrawal syndrome, although qualitatively similar to that of morphine, differs in that the onset is slower, the course is more prolonged, and the symptoms are less severe.

Some data also indicate that methadone acts as an antagonist at the N-methyl-D-aspartate (NMDA) receptor. The contribution of NMDA receptor antagonism to methadone's efficacy is unknown. Other NMDA receptor antagonists have been shown to produce neurotoxic effects in animals.

12.3 Pharmacokinetics

Absorption: Following oral administration the bioavailability of methadone ranges between 36 to 100% and peak plasma concentrations are reached within 1 to 7.5 hours. Dose proportionality of methadone pharmacokinetics is not known. However, after administration of daily oral doses ranging from 10 to 225 mg, the steady-state plasma concentrations ranged between 65 to 630 ng/mL and the peak concentrations ranged between 124 to 1255 ng/mL. Effect of food on the bioavailability of methadone has not been evaluated.

Distribution: Methadone is a lipophilic drug and the steady-state volume of distribution ranges between 1.0 to 8.0 L/kg. In plasma, methadone is predominantly bound to α1-acid glycoprotein (85% to 90%). Methadone is secreted in saliva, breast milk, amniotic fluid and urine.

Metabolism: Methadone is primarily metabolized by N-demethylation to an inactive metabolite, 2-ethylidene-1,5-dimethyl-3,3-diphenylpyrrolidine (EDDP). Cytochrome P450 enzymes, primarily CYP3A4, CYP2B6, and CYP2C19 and to a lesser extent CYP2C9 and CYP2D6, are responsible for conversion of methadone to EDDP and other inactive metabolites, which are excreted mainly in the urine. Methadone appears to be a substrate for P-glycoprotein but its pharmacokinetics do not appear to be significantly altered in case of P-glycoprotein polymorphism or inhibition.

Excretion: The elimination of methadone is mediated by extensive biotransformation, followed by renal and fecal excretion. The elimination half-life of methadone is approximately 24 hours. The apparent plasma clearance of methadone ranged between 1.4 and 126 L/h, and the terminal half-life (T_{1/2}) was highly variable and ranged between 8 to 59 hours in different studies. Methadone is a basic (pKa=9.2) compound and the pH of the urinary tract after its disposition in plasma. Also, since methadone is lipophilic, it has been known to persist in the liver and other tissues. The slow release from the liver and other tissues may prolong the duration of methadone action despite low plasma concentrations.

Drug Interactions: Cytochrome P450 Interactions: Methadone undergoes hepatic N-demethylation by CYP3A4, CYP2B6, and CYP2C19, and to a lesser extent by CYP2C9 and CYP2D6. Co-administration of methadone with CYP inducers may result in more rapid metabolism and potential for decreased effects of methadone, whereas administration with CYP inhibitors may reduce metabolism and potentiate methadone's effects. Although antiretroviral drugs such as efavirenz, nelfinavir, nevirapine, ritonavir, lopinavir+ritonavir combination are known to inhibit some CYPs, they are shown to reduce the plasma levels of methadone, possibly due to CYP induction activity [see *Drug Interactions* (7.2)]. Therefore, drugs administered concomitantly with methadone should be evaluated for interaction potential; clinicians are advised to evaluate individual response to drug therapy.

Cytochrome P450 Inducers: The following drug interactions were reported following coadministration of methadone with known inducers of cytochrome P450 enzymes:

Rifampin: In patients well-stabilized on methadone, concomitant administration of rifampin resulted in a marked reduction in serum methadone levels and a concurrent appearance of withdrawal symptoms.

Phenytoin: In a pharmacokinetic study with patients on methadone maintenance therapy, phenytoin administration (250 mg twice daily initially for 1 day followed by 300 mg daily for 3 to 4 days) resulted in an approximately 50% reduction in methadone exposure and withdrawal symptoms occurred concurrently. Upon discontinuation of phenytoin, the incidence of withdrawal symptoms decreased and methadone exposure increased to a level comparable to that prior to phenytoin administration.

St. John's Wort, Phenobarbital, Carbamazepine: Administration of methadone with other CYP3A4 inducers may result in withdrawal symptoms.

Cytochrome P450 Inhibitors: Since the metabolism of methadone is mediated primarily by CYP3A4 isozyme, coadministration of drugs that inhibit CYP3A4 activity may cause decreased clearance of methadone.

Variconazole: Repeat dose administration of oral variconazole (400 mg every 12 hours for 1 day, then 200 mg every 12 hours for 4 days) increased the peak plasma concentration (C_{max}) and AUC (C₀₋₂₄) of methadone by 21% and 47%, respectively, in subjects receiving a methadone maintenance dose (30 to 100 mg daily). The C_{max} and AUC of (S)-methadone increased by 65% and 103%, respectively. Increased plasma concentrations of methadone have been associated with toxicity including QT prolongation. Frequent monitoring for adverse events and toxicity related to methadone is recommended during coadministration. Dose reduction of methadone may be needed [see *Drug Interactions* (7.2)].

Antiretroviral Drugs: Although antiretroviral drugs such as efavirenz, nelfinavir, nevirapine, ritonavir, telaprevir, lopinavir+ritonavir combination are known to inhibit some CYPs, they are shown to reduce the plasma levels of methadone, possibly due to CYP induction activity [see *Drug Interactions* (7.2)].

Abacavir, amprevnavir, darunavir+ritonavir, efavirenz, nelfinavir, nevirapine, ritonavir, telaprevir, lopinavir+ritonavir, saquinavir+ritonavir, tipranavir+ritonavir combination: Coadministration of these antiretroviral agents resulted in increased clearance or decreased plasma levels of methadone [see *Drug Interactions* (7.2)].

Didanosine and Stavudine: Methadone decreased the AUC and peak levels for didanosine and stavudine, with a more significant decrease for didanosine. Methadone disposition was not substantially altered [see *Drug Interactions* (7.2)].

Zidovudine: Methadone increased the AUC of zidovudine which could result in toxic effects [see *Drug Interactions* (7.2)].

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis: The results of carcinogenicity assessment in B6C2F1 mice and Fischer 344 rats following dietary administration of two doses of methadone HCl have been published. Mice consumed 15 mg/kg/ day or 60 mg/kg/day for two years. Rats consumed 15 mg/kg/day or 60 mg/kg/day for two years. These doses were approximately 0.6 and 2.5 times a human daily oral dose of 120 mg/day on a body surface area basis (mg/m²). There was a significant increase in pituitary adenomas in female mice treated with 15 mg/kg/day but not with 60 mg/kg/day. Under the conditions of the assay, there was no clear evidence for a treatment-related increase in the incidence of neoplasms in male rats. Due to decreased food consumption in males at the high dose, male rats consumed 16 mg/kg/day and 28 mg/kg/day of methadone for two years. These doses were approximately 1.3 and 2.3 times a human daily oral dose of 120 mg/day, based on body surface area comparison. In contrast, female rats consumed 16 mg/kg/day and 28 mg/kg/day for two years. These doses were approximately 3.7 and 7.1 times a human daily oral dose of 120 mg/day, based on body surface area comparison. Under the conditions of the assay, there was no clear evidence for a treatment-related increase in the incidence of neoplasms in either male or female rats.

Mutagenesis: There are several published reports on the potential genetic toxicity of methadone. Methadone tested positive in the *in vivo* mouse dominant lethal assay and the *in vivo* mammalian spermatogonial chromosome aberration test. Additionally, methadone tested positive in the *E. coli* DNA repair system and *Neurospora crassa* and mouse lymphoma forward mutation assays. In contrast, methadone tested negative in tests for chromosome breakage and disjunction and sex-linked recessive lethal gene mutations in germ cells of *Drosophila* using feeding and injection procedures.

Fertility: Published animal studies show that methadone treatment of males can alter reproductive function. Methadone produces a significant regression of sex accessory organs and testes of male mice and rats.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 Storage and Handling

Methadone is a controlled substance. Like fentanyl, morphine, oxycodone, hydromorphone, and oxycodone, methadone is controlled under Schedule II of the Federal Controlled Substances Act. Methadone may be targeted for theft and diversion by criminals [see *Warnings and Precautions* (5.1)].

Dispense in a tight, light-resistant container as defined in the USP/NF.

Store at 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].

16.2 How Supplied

Methadone Hydrochloride Oral Solution USP

Methadone Hydrochloride Oral Solution USP 5 and 10 mgper 5 mL is an orange-colored, citrus-flavored solution available in two strengths as follows:

5 mg per 5 mL Oral Solution
NDC 0054-3555-63: Bottles of 500 mL

10 mg per 5 mL Oral Solution
NDC 0054-3556-63: Bottles of 500 mL

DEA order form required.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

Addition, Abuse, and Misuse: Inform patients that the use of methadone, even when taken as recommended, can result in addiction, abuse, and misuse, which can lead to overdose or death [see *Warnings and Precautions* (5.1)]. Instruct patients not to share methadone with others and to take steps to protect methadone from theft or misuse.

Life-threatening Respiratory Depression: Inform patients of the risk of life-threatening respiratory depression, including information that the risk is greatest when starting methadone or when the dose is increased, and that it can occur even at recommended doses [see *Warnings and Precautions* (5.2)]. Advise patients how to recognize respiratory depression and to seek medical attention if breathing difficulties develop.

Accidental Ingestion: Inform patients that accidental ingestion, especially in children, may result in respiratory depression or death [see *Warnings and Precautions* (5.2)]. Instruct patients to take steps to store methadone securely and to dispose of unused methadone by flushing the tablets down the toilet.

Symptoms of Arrhythmia: Instruct patients to seek medical attention immediately if they experience symptoms suggestive of an arrhythmia (such as palpitations, near syncope, or syncope) when taking methadone.

Neonatal Opioid Withdrawal Syndrome: Inform female patients of reproductive potential that prolonged use of methadone during pregnancy can result in neonatal opioid withdrawal syndrome, which may be life-threatening if not recognized and treated [see *Warnings and Precautions* (5.4)].

Interactions with Alcohol and other CNS Depressants: Inform patients that potentially serious additive effects may occur if methadone is used with alcohol or other CNS depressants, and not to use such drugs unless supervised by a health care provider.

Important Administration Instructions: Instruct patients how to properly take methadone, including the following:

- Use methadone exactly as prescribed to reduce the risk of life-threatening adverse reactions (e.g., respiratory depression)
- Do not discontinue methadone without first discussing the need for a tapering regimen with the prescriber

Hypotension: Inform patients that methadone may cause orthostatic hypotension and syncope. Instruct patients how to recognize symptoms of low blood pressure and how to reduce the risk of serious consequences should hypotension occur (e.g., sit or lie down, carefully rise from a sitting or lying position).

Driving or Operating Heavy Machinery: Inform patients that methadone may impair the ability to perform potentially hazardous activities such as driving a car or operating heavy machinery. Advise patients not to perform such tasks until they know how they will react to the medication.

Constipation: Advise patients of the potential for severe constipation, including management instructions and when to seek medical attention.

Anaphylaxis: Inform patients that anaphylaxis has been reported with ingredients contained in methadone. Advise patients how to recognize such a reaction and when to seek medical attention.

Breastfeeding: Instruct nursing mothers using methadone to watch for signs of methadone toxicity in their infants, which include increased sleepiness (more than usual), difficulty breastfeeding, breathing difficulties, or limpness. Instruct nursing mothers to talk to the baby's healthcare provider immediately if they notice these signs. If they cannot reach the healthcare provider right away, instruct them to take the baby to the emergency room or call 911 (or local emergency services).

Disposal of Unused Methadone: Advise patients to flush the unused tablets down the toilet when methadone is no longer needed.

Roxane Laboratories, Inc.
Columbus, Ohio 43216

4056301/06

Boehringer Ingelheim
Roxane Laboratories

Revised April 2014
© RLI, 2014